

PROBLEMS IN ELEMENTARY NUMBER THEORY

Hojoo Lee

Version 050722

God does arithmetic. **C. F. Gauss**

1, -24, 252, -1472, 4830, -6048, -16744, 84480, -113643, -115920,
534612, -370944, -577738, 401856, 1217160, 987136, -6905934, 2727432,
10661420, -7109760, -4219488, -12830688, 18643272, 21288960, -25499225,
13865712, -73279080, 24647168, ...

CONTENTS

1.	Introduction	3
2.	Notations and Abbreviations	4
3.	Divisibility Theory I	5
4.	Divisibility Theory II	12
5.	Arithmetic in Z_n	16
	Primitive Roots	16
	Quadratic Residues	17
	Congruences	17
6.	Primes and Composite Numbers	20
	Composite Numbers	20
	Prime Numbers	20
7.	Rational and Irrational Numbers	24
	Rational Numbers	24
	Irrational Numbers	25
8.	Diophantine Equations I	29
9.	Diophantine Equations II	34
10.	Functions in Number Theory	37
	Floor Function and Fractional Part Function	37
	Euler phi Function	39
	Divisor Functions	39
	More Functions	40
	Functional Equations	41
11.	Polynomials	44
12.	Sequences of Integers	46
	Linear Recurrences	46
	Recursive Sequences	47
	More Sequences	51
13.	Combinatorial Number Theory	54
14.	Additive Number Theory	61
15.	The Geometry of Numbers	66
16.	Miscellaneous Problems	67
17.	Sources	71
18.	References	94

1. INTRODUCTION

The heart of Mathematics is its problems. Paul Halmos

1. Introduction *Number Theory* is a beautiful branch of Mathematics. The purpose of this book is to present a collection of interesting questions in *Number Theory*. Many of the problems are mathematical competition problems all over the world including IMO, APMO, APMC, and Putnam, etc. The book is available at

<http://my.netian.com/~ideahitme/orange.html>

2. How You Can Help This is an **unfinished** manuscript. I would greatly appreciate hearing about any errors in the book, even minor ones. I also would like to hear about

- a) *challenging* problems in *elementary number theory*,
- b) *interesting* problems concerned with the *history of number theory*,
- c) *beautiful* results that are *easily* stated, and
- d) *remarks* on the problems in the book.

You can send all comments to the author at **ultrametric@gmail.com**

3. Acknowledgments The author is very grateful to *Orlando Doehring*, who provided old IMO short-listed problems. The author also wish to thank *Arne Smeets*, *Ha Duy Hung*, *Tom Verhoeff* and *Tran Nam Dung* for their nice problem proposals and comments.

2. NOTATIONS AND ABBREVIATIONS

Notations

\mathbf{Z} is the set of integers

\mathbf{N} is the set of positive integers

\mathbf{N}_0 is the set of nonnegative integers

\mathbb{Q} is the set of rational numbers

$m|n$ n is a multiple of m .

$\sum_{d|n} f(d) = \sum_{d \in D(n)} f(d)$ ($D(n) = \{d \in \mathbf{N} : d|n\}$)

$[x]$ the greatest integer less than or equal to x

$\{x\}$ the fractional part of x ($\{x\} = x - [x]$)

$\pi(x)$ the number of primes p with $p \leq x$

$\phi(n)$ the number of positive integers less than n that are relatively prime to n

$\sigma(n)$ the sum of positive divisors of n

$d(n)$ the number of positive divisors of n

τ Ramanujan's tau function

Abbreviations

AIME American Invitational Mathematics Examination

APMO Asian Pacific Mathematics Olympiads

IMO International Mathematical Olympiads

CRUX Crux Mathematicorum (with Mathematical Mayhem)

3. DIVISIBILITY THEORY I

Why are numbers beautiful? It's like asking why is Beethoven's Ninth Symphony beautiful. If you don't see why, someone can't tell you. I know numbers are beautiful. If they aren't beautiful, nothing is. Paul Erdős

A 1. (Kiran S. Kedlaya) Show that if x, y, z are positive integers, then $(xy + 1)(yz + 1)(zx + 1)$ is a perfect square if and only if $xy + 1$, $yz + 1$, $zx + 1$ are all perfect squares.

A 2. Find infinitely many triples (a, b, c) of positive integers such that a, b, c are in arithmetic progression and such that $ab + 1$, $bc + 1$, and $ca + 1$ are perfect squares.

A 3. Let a and b be positive integers such that $ab + 1$ divides $a^2 + b^2$. Show that

$$\frac{a^2 + b^2}{ab + 1}$$

is the square of an integer.

A 4. (Shailesh Shirali) If a, b, c are positive integers such that

$$0 < a^2 + b^2 - abc \leq c,$$

show that $a^2 + b^2 - abc$ is a perfect square.¹

A 5. Let x and y be positive integers such that xy divides $x^2 + y^2 + 1$. Show that

$$\frac{x^2 + y^2 + 1}{xy} = 3.$$

A 6. (R. K. Guy and R. J. Nowakowski) (i) Find infinitely many pairs of integers a and b with $1 < a < b$, so that ab exactly divides $a^2 + b^2 - 1$. (ii) With a and b as in (i), what are the possible values of

$$\frac{a^2 + b^2 - 1}{ab}.$$

A 7. Let n be a positive integer such that $2 + 2\sqrt{28n^2 + 1}$ is an integer. Show that $2 + 2\sqrt{28n^2 + 1}$ is the square of an integer.

A 8. The integers a and b have the property that for every nonnegative integer n the number of $2^n a + b$ is the square of an integer. Show that $a = 0$.

A 9. Prove that among any ten consecutive positive integers at least one is relatively prime to the product of the others.

¹This is a generalization of **A3**! Indeed, $a^2 + b^2 - abc = c$ implies that $\frac{a^2 + b^2}{ab + 1} = c \in \mathbf{N}$.

A 10. Let n be a positive integer with $n \geq 3$. Show that

$$n^{n^{n^n}} - n^{n^n}$$

is divisible by 1989.

A 11. Let a, b, c, d be integers. Show that the product

$$(a - b)(a - c)(a - d)(b - c)(b - d)(c - d)$$

is divisible by 12.²

A 12. Let k, m , and n be natural numbers such that $m + k + 1$ is a prime greater than $n + 1$. Let $c_s = s(s + 1)$. Prove that the product

$$(c_{m+1} - c_k)(c_{m+2} - c_k) \cdots (c_{m+n} - c_k)$$

is divisible by the product $c_1 c_2 \cdots c_n$.

A 13. Show that for all prime numbers p ,

$$Q(p) = \prod_{k=1}^{p-1} k^{2k-p-1}$$

is an integer.

A 14. Let n be an integer with $n \geq 2$. Show that n does not divide $2^n - 1$.

A 15. Suppose that $k \geq 2$ and $n_1, n_2, \dots, n_k \geq 1$ be natural numbers having the property

$$n_2 \mid 2^{n_1} - 1, n_3 \mid 2^{n_2} - 1, \dots, n_k \mid 2^{n_{k-1}} - 1, n_1 \mid 2^{n_k} - 1.$$

Show that $n_1 = n_2 = \cdots = n_k = 1$.

A 16. Determine if there exists a positive integer n such that n has exactly 2000 prime divisors and $2^n + 1$ is divisible by n .

A 17. Let m and n be natural numbers such that

$$A = \frac{(m+3)^n + 1}{3m}.$$

is an integer. Prove that A is odd.

A 18. Let m and n be natural numbers and let $mn + 1$ be divisible by 24. Show that $m + n$ is divisible by 24.

A 19. Let $f(x) = x^3 + 17$. Prove that for each natural number $n \geq 2$, there is a natural number x for which $f(x)$ is divisible by 3^n but not 3^{n+1} .

A 20. Determine all positive integers n for which there exists an integer m so that $2^n - 1$ divides $m^2 + 9$.

A 21. Let n be a positive integer. Show that the product of n consecutive integers is divisible by $n!$

²There is a strong generalization. See **J1**

A 22. Prove that the number

$$\sum_{k=0}^n (2n+1) \binom{2n+1}{2k+1} 2^{3k}$$

is not divisible by 5 for any integer $n \geq 0$.

A 23. (Wolstenholme's Theorem) Prove that if

$$1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{p-1}$$

is expressed as a fraction, where $p \geq 5$ is a prime, then p^2 divides the numerator.

A 24. If p is a prime number greater than 3 and $k = \lfloor \frac{2p}{3} \rfloor$. Prove that

$$\binom{p}{1} + \binom{p}{2} + \cdots + \binom{p}{k}$$

is divisible by p^2 .

A 25. Show that $\binom{2n}{n} \mid \text{lcm}[1, 2, \dots, 2n]$ for all positive integers n .

A 26. Let m and n be arbitrary non-negative integers. Prove that

$$\frac{(2m)!(2n)!}{m!n!(m+n)!}$$

is an integer. ($0! = 1$).

A 27. Show that the coefficients of a binomial expansion $(a+b)^n$ where n is a positive integer, are all odd, if and only if n is of the form $2^k - 1$ for some positive integer k .

A 28. Prove that the expression

$$\frac{\text{gcd}(m,n)}{n} \binom{n}{m}$$

is an integer for all pairs of positive integers (m, n) with $n \geq m \geq 1$.

A 29. For which positive integers k , is it true that there are infinitely many pairs of positive integers (m, n) such that

$$\frac{(m+n-k)!}{m!n!}$$

is an integer?

A 30. Show that if $n \geq 6$ is composite, then n divides $(n-1)!$.

A 31. Show that there exist infinitely many positive integers n such that $n^2 + 1$ divides $n!$.

A 32. Let p and q be natural numbers such that

$$\frac{p}{q} = 1 - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \cdots - \frac{1}{1318} + \frac{1}{1319}.$$

Prove that p is divisible by 1979.

A 33. Let $b > 1$, a and n be positive integers such that $b^n - 1$ divides a . Show that in base b , the number a has at least n non-zero digits.

A 34. Let p_1, p_2, \dots, p_n be distinct primes greater than 3. Show that

$$2^{p_1 p_2 \cdots p_n} + 1$$

has at least 4^n divisors.

A 35. Let $p \geq 5$ be a prime number. Prove that there exists an integer a with $1 \leq a \leq p - 2$ such that neither $a^{p-1} - 1$ nor $(a + 1)^{p-1} - 1$ is divisible by p^2 .

A 36. Let n and q be integers with $n \geq 5$, $2 \leq q \leq n$. Prove that $q - 1$ divides $\left\lfloor \frac{(n-1)!}{q} \right\rfloor$.

A 37. If n is a natural number, prove that the number $(n+1)(n+2) \cdots (n+10)$ is not a perfect square.

A 38. Let p be a prime with $p > 5$, and let $S = \{p - n^2 \mid n \in \mathbf{N}, n^2 < p\}$. Prove that S contains two elements a and b such that $a \mid b$ and $1 < a < b$.

A 39. Let n be a positive integer. Prove that the following two statements are equivalent.

- n is not divisible by 4
- There exist $a, b \in \mathbf{Z}$ such that $a^2 + b^2 + 1$ is divisible by n .

A 40. Determine the greatest common divisor of the elements of the set

$$\{n^{13} - n \mid n \in \mathbf{Z}\}.$$

A 41. Show that there are infinitely many composite n such that $3^{n-1} - 2^{n-1}$ is divisible by n .

A 42. Suppose that $2^n + 1$ is an odd prime for some positive integer n . Show that n must be a power of 2.

A 43. Suppose that p is a prime number and is greater than 3. Prove that $7^p - 6^p - 1$ is divisible by 43.

A 44. Suppose that $4^n + 2^n + 1$ is prime for some positive integer n . Show that n must be a power of 3.

A 45. Let b , m , and n be positive integers $b > 1$ and m and n are different. Suppose that $b^m - 1$ and $b^n - 1$ have the same prime divisors. Show that $b + 1$ must be a power of 2.

A 46. Let a and b be integers. Show that a and b have the same parity if and only if there exist integers c and d such that $a^2 + b^2 + c^2 + 1 = d^2$.

A 47. Let n be a positive integer with $n > 1$. Prove that

$$\frac{1}{2} + \cdots + \frac{1}{n}$$

is not an integer.

A 48. Let n be a positive integer. Prove that

$$\frac{1}{3} + \cdots + \frac{1}{2n+1}$$

is not an integer.

A 49. Prove that there is no positive integer n such that, for $k = 1, 2, \dots, 9$, the leftmost digit (in decimal notation) of $(n+k)!$ equals k .

A 50. Show that every integer $k > 1$ has a multiple less than k^4 whose decimal expansion has at most four distinct digits.

A 51. Let a, b, c and d be odd integers such that $0 < a < b < c < d$ and $ad = bc$. Prove that if $a + d = 2^k$ and $b + c = 2^m$ for some integers k and m , then $a = 1$.

A 52. Let d be any positive integer not equal to 2, 5, or 13. Show that one can find distinct a and b in the set $\{2, 5, 13, d\}$ such that $ab - 1$ is not a perfect square.

A 53. Suppose that x, y , and z are positive integers with $xy = z^2 + 1$. Prove that there exist integers a, b, c , and d such that $x = a^2 + b^2$, $y = c^2 + d^2$, and $z = ac + bd$.

A 54. A natural number n is said to have the property P , if whenever n divides $a^n - 1$ for some integer a , n^2 also necessarily divides $a^n - 1$.

- (a) Show that every prime number n has the property P .
 (b) Show that there are infinitely many composite numbers n that possess the property P .

A 55. Show that for every natural number n the product

$$\left(4 - \frac{2}{1}\right) \left(4 - \frac{2}{2}\right) \left(4 - \frac{2}{3}\right) \cdots \left(4 - \frac{2}{n}\right)$$

is an integer.

A 56. Let a, b , and c be integers such that $a + b + c$ divides $a^2 + b^2 + c^2$. Prove that there are infinitely many positive integers n such that $a + b + c$ divides $a^n + b^n + c^n$.

A 57. Prove that for every $n \in \mathbf{N}$ the following proposition holds : The number 7 is a divisor of $3^n + n^3$ if and only if 7 is a divisor of $3^n n^3 + 1$.

A 58. Let $k \geq 14$ be an integer, and let p_k be the largest prime number which is strictly less than k . You may assume that $p_k \geq 3k/4$. Let n be a composite integer. Prove that

- (a) if $n = 2p_k$, then n does not divide $(n - k)!$
 (b) if $n > 2p_k$, then n divides $(n - k)!$.

A 59. Suppose that n has (at least) two essentially distinct representations as a sum of two squares. Specifically, let $n = s^2 + t^2 = u^2 + v^2$, where $s \geq t \geq 0$, $u \geq v \geq 0$, and $s > u$. Show that $\gcd(su - tv, n)$ is a proper divisor of n .

A 60. Prove that there exist an infinite number of ordered pairs (a, b) of integers such that for every positive integer t , the number $at + b$ is a triangular number if and only if t is a triangular number³.

A 61. For any positive integer $n > 1$, let $p(n)$ be the greatest prime divisor of n . Prove that there are infinitely many positive integers n with

$$p(n) < p(n + 1) < p(n + 2).$$

A 62. Let $p(n)$ be the greatest odd divisor of n . Prove that

$$\frac{1}{2^n} \sum_{k=1}^{2^n} \frac{p(k)}{k} > \frac{2}{3}.$$

A 63. There is a large pile of cards. On each card one of the numbers $1, 2, \dots, n$ is written. It is known that the sum of all numbers of all the cards is equal to $k \cdot n!$ for some integer k . Prove that it is possible to arrange cards into k stacks so that the sum of numbers written on the cards in each stack is equal to $n!$.

A 64. The last digit of the number $x^2 + xy + y^2$ is zero (where x and y are positive integers). Prove that two last digits of this numbers are zeros.

A 65. Clara computed the product of the first n positive integers and Valerid computed the product of the first m even positive integers, where $m \geq 2$. They got the same answer. Prove that one of them had made a mistake.

A 66. (Four Number Theorem) Let a, b, c , and d be positive integers such that $ab = cd$. Show that there exists positive integers p, q, r , and s such that

$$a = pq, \quad b = rs, \quad c = pt, \quad \text{and} \quad d = su.$$

A 67. Prove that $\binom{2n}{n}$ is divisible by $n + 1$.

A 68. Suppose that a_1, \dots, a_r are positive integers. Show that $\text{lcm}[a_1, \dots, a_r] = a_1 \cdots a_r (a_1, a_2)^{-1} \cdots (a_{r-1}, a_r)^{-1} (a_1, a_2, a_3)(a_1, a_2, a_3) \cdots (a_1, a_2, \dots, a_r)^{(-1)^{r+1}}$.

A 69. Prove that if the odd prime p divides $a^b - 1$, where a and b are positive integers, then p appears to the same power in the prime factorization of $b(a^d - 1)$, where d is the greatest common divisor of b and $p - 1$.

³The triangular numbers are the $t_n = n(n + 1)/2$ with $n \in \{0, 1, 2, \dots\}$.

A 70. Suppose that $m = nq$, where n and q are positive integers. Prove that the sum of binomial coefficients

$$\sum_{k=0}^{n-1} \binom{(n,k)q}{(n,k)}$$

is divisible by m , where (x, y) denotes the greatest common divisor of x and y .

4. DIVISIBILITY THEORY II

Number theorists are like lotus-eaters - having tasted this food they can never give it up. Leopold Kronecker

B 1. Determine all integers $n > 1$ such that

$$\frac{2^n + 1}{n^2}$$

is an integer.

B 2. Determine all pairs (n, p) of nonnegative integers such that

- p is a prime,
- $n < 2p$, and
- $(p - 1)^n + 1$ is divisible by n^{p-1} .

B 3. Determine all pairs (n, p) of positive integers such that

- p is a prime, $n > 1$, and
- $(p - 1)^n + 1$ is divisible by n^{p-1} .⁴

B 4. Find an integer n , where $100 \leq n \leq 1997$, such that

$$\frac{2^n + 2}{n}$$

is also an integer.

B 5. Find all triples (a, b, c) of positive integers such that $2^c - 1$ divides $2^a + 2^b + 1$.

B 6. Find all integers a, b, c with $1 < a < b < c$ such that

$$(a - 1)(b - 1)(c - 1) \quad \text{is a divisor of} \quad abc - 1.$$

B 7. Find all positive integers, representable uniquely as

$$\frac{x^2 + y}{xy + 1},$$

where x and y are positive integers.

B 8. Determine all ordered pairs (m, n) of positive integers such that

$$\frac{n^3 + 1}{mn - 1}$$

is an integer.

⁴The answer is $(n, p) = (2, 2), (3, 3)$. Note that this problem is a very nice generalization of the above two IMO problems B1 and B2 !

B 9. Determine all pairs of integers (a, b) such that

$$\frac{a^2}{2a^2b - b^3 + 1}$$

is a positive integer.

B 10. Find all pairs of positive integers $m, n \geq 3$ for which there exist infinitely many positive integers a such that

$$\frac{a^m + a - 1}{a^n + a^2 - 1}$$

is itself an integer.

B 11. Determine all triples of positive integers (a, m, n) such that $a^m + 1$ divides $(a + 1)^n$.

B 12. Which integers are represented by $\frac{(x+y+z)^2}{xyz}$ where $x, y,$ and z are positive integers?

B 13. Find all $n \in \mathbf{N}$ such that $[\sqrt{n}] \mid n$.

B 14. Determine all $n \in \mathbf{N}$ for which (i) n is not the square of any integer, and (ii) $[\sqrt{n}]^3$ divides n^2 .

B 15. Find all $n \in \mathbf{N}$ such that $2^{n-1} \mid n!$.

B 16. Find all positive integers (x, n) such that $x^n + 2^n + 1$ is a divisor of $x^{n+1} + 2^{n+1} + 1$.

B 17. Find all positive integers n such that $3^n - 1$ is divisible by 2^n .

B 18. Find all positive integers n such that $9^n - 1$ is divisible by 7^n .

B 19. Determine all pairs (a, b) of integers for which $a^2 + b^2 + 3$ is divisible by ab .

B 20. Determine all pairs (x, y) of positive integers with $y \mid x^2 + 1$ and $x \mid y^3 + 1$.

B 21. Determine all pairs (a, b) of positive integers such that $ab^2 + b + 7$ divides $a^2b + a + b$.

B 22. Let a and b be positive integers. When $a^2 + b^2$ is divided by $a + b$, the quotient is q and the remainder is r . Find all pairs (a, b) such that $q^2 + r = 1977$.

B 23. Find the largest positive integer n such that n is divisible by all the positive integers less than $n^{1/3}$.

B 24. Find all $n \in \mathbf{N}$ such that $3^n - n$ is divisible by 17.

B 25. Suppose that a and b are natural numbers such that

$$p = \frac{4}{b} \sqrt{\frac{2a-b}{2a+b}}$$

is a prime number. What is the maximum possible value of p ?

B 26. Find all positive integers n that have exactly 16 positive integral divisors d_1, d_2, \dots, d_{16} such that $1 = d_1 < d_2 < \dots < d_{16} = n$, $d_6 = 18$, and $d_9 - d_8 = 17$.

B 27. Suppose that n is a positive integer and let

$$d_1 < d_2 < d_3 < d_4$$

be the four smallest positive integer divisors of n . Find all integers n such that

$$n = d_1^2 + d_2^2 + d_3^2 + d_4^2.$$

B 28. Let $1 = d_1 < d_2 < \dots < d_k = n$ be all different divisors of positive integer n written in ascending order. Determine all n such that

$$d_7^2 + d_{10}^2 = \left(\frac{n}{d_{22}}\right)^2.$$

B 29. Let $n \geq 2$ be a positive integer, with divisors

$$1 = d_1 < d_2 < \dots < d_k = n.$$

Prove that

$$d_1 d_2 + d_2 d_3 + \dots + d_{k-1} d_k$$

is always less than n^2 , and determine when it is a divisor of n^2 .

B 30. Find all positive integers n such that (a) n has exactly 6 positive divisors $1 < d_1 < d_2 < d_3 < d_4 < n$, and (b) $1 + n = 5(d_1 + d_2 + d_3 + d_4)$.

B 31. Find all composite numbers n , having the property : each divisor d of n ($d \neq 1, n$) satisfies inequalities $n - 20 \leq d \leq n - 12$.

B 32. Determine all three-digit numbers N having the property that N is divisible by 11, and $\frac{N}{11}$ is equal to the sum of the squares of the digits of N .

B 33. When 4444^{4444} is written in decimal notation, the sum of its digits is A . Let B be the sum of the digits of A . Find the sum of the digits of B . (A and B are written in decimal notation.)

B 34. A wobbly number is a positive integer whose digits in base 10 are alternatively non-zero and zero the units digit being non-zero. Determine all positive integers which do not divide any wobbly number.

B 35. Find the smallest positive integer n such that

- (i) n has exactly 144 distinct positive divisors, and
- (ii) there are ten consecutive integers among the positive divisors of n .

B 36. Determine the least possible value of the natural number n such that $n!$ ends in exactly 1987 zeros.

B 37. Find four positive integers, each not exceeding 70000 and each having more than 100 divisors.

B 38. For each integer $n > 1$, let $p(n)$ denote the largest prime factor of n . Determine all triples (x, y, z) of distinct positive integers satisfying

- (i) x, y, z are in arithmetic progression, and
(ii) $p(xyz) \leq 3$.

B 39. Find all positive integers a and b such that

$$\frac{a^2 + b}{b^2 - a} \quad \text{and} \quad \frac{b^2 + a}{a^2 - b}$$

are both integers.

B 40. For each positive integer n , write the sum $\sum_{m=1}^n 1/m$ in the form p_n/q_n , where p_n and q_n are relatively prime positive integers. Determine all n such that 5 does not divide q_n .

B 41. Find all natural numbers n such that the number $n(n+1)(n+2)(n+3)$ has exactly three prime divisors.

B 42. Prove that there exist infinitely many pairs (a, b) of relatively prime positive integers such that

$$\frac{a^2 - 5}{b} \quad \text{and} \quad \frac{b^2 - 5}{a}$$

are both positive integers.

B 43. Determine all triples (l, m, n) of distinct positive integers satisfying

$$\gcd(l, m)^2 = l + m, \quad \gcd(m, n)^2 = m + n, \quad \text{and} \quad \gcd(n, l)^2 = n + l.$$

B 44. What is the greatest common divisor of the set of numbers

$$\{16^n + 10n - 1 \mid n = 1, 2, \dots\}?$$

B 45. (I. Selishev) Does there exist a 4-digit integer (in decimal form) such that no replacement of three of its digits by another three gives a multiple of 1992 ?

B 46. What is the smallest positive integer that consists of the ten digits 0 through 9, each used just once, and is divisible by each of the digits 2 through 9 ?

B 47. Find the smallest positive integer n which makes

$$2^{1989} \mid m^n - 1$$

for all odd positive integer m greater than 1.

B 48. Determine the highest power of 1980 which divides

$$\frac{(1980n)!}{(n!)^{1980}}.$$

5. ARITHMETIC IN Z_n

Mathematics is the queen of the sciences and number theory is the queen of Mathematics. Johann Carl Friedrich Gauss

5.1. Primitive Roots.

C 1. Let n be a positive integer. Show that there are infinitely many primes p such that the smallest positive primitive root of p is greater than n .

C 2. Let p be a prime with $p > 4 \left(\frac{p-1}{\phi(p-1)} \right)^2 2^{2k}$, where k denotes the number of distinct prime divisors of $p-1$, and let M be an integer. Prove that the set of integers $M+1, M+2, \dots, M+2 \left\lceil \frac{p-1}{\phi(p-1)} 2^k \sqrt{p} \right\rceil - 1$ contains a primitive root to modulus p .

C 3. Show that for each odd prime p , there is an integer g such that $1 < g < p$ and g is a primitive root modulo p^n for every positive integer n .

C 4. Let g be a Fibonacci primitive root (mod p). i.e. g is a primitive root (mod p) satisfying $g^2 \equiv g+1 \pmod{p}$. Prove that

- (a) Prove that $g-1$ is also a primitive root (mod p).
- (b) If $p = 4k+3$, then $(g-1)^{2k+3} \equiv g-2 \pmod{p}$ and deduce that $g-2$ is also a primitive root (mod p).

C 5. Let p be an odd prime. If $g_1, \dots, g_{\phi(p-1)}$ are the primitive roots mod p in the range $1 < g \leq p-1$, prove that

$$\sum_{i=1}^{\phi(p-1)} g_i \equiv \mu(p-1) \pmod{p}.$$

C 6. Suppose that m does not have a primitive root. Show that

$$a^{\frac{\phi(m)}{2}} \equiv -1 \pmod{m}$$

for every a relatively prime m .

C 7. Suppose that $p > 3$ is prime. Prove that the products of the primitive roots of p between 1 and $p-1$ is congruent to 1 modulo p .

C 8. Let p be a prime. Let g be a primitive root of modulo p . Prove that there is no k such that $g^{k+2} \equiv g^{k+1} + 1 \equiv g^k + 2 \pmod{p}$.

5.2. Quadratic Residues.

C 9. Find all positive integers n that are quadratic residues modulo all primes greater than n .

C 10. The positive integers a and b are such that the numbers $15a + 16b$ and $16a - 15b$ are both squares of positive integers. What is the least possible value that can be taken on by the smaller of these two squares?

C 11. Let p be an odd prime number. Show that the smallest positive quadratic nonresidue of p is smaller than $\sqrt{p} + 1$.

C 12. Let M be an integer, and let p be a prime with $p > 25$. Show that the sequence $M, M + 1, \dots, M + 3[\sqrt{p}] - 1$ contains a quadratic non-residue to modulus p .

C 13. Let p be an odd prime and let Z_p denote (the field of) integers modulo p . How many elements are in the set

$$\{x^2 : x \in Z_p\} \cap \{y^2 + 1 : y \in Z_p\}?$$

C 14. Let a, b, c be integers and let p be an odd prime with

$$p \nmid a \text{ and } p \nmid b^2 - 4ac.$$

Show that

$$\sum_{k=1}^p \left(\frac{ak^2 + bk + c}{p} \right) = - \left(\frac{a}{p} \right).$$

5.3. Congruences.

C 15. If p is an odd prime, prove that

$$\binom{k}{p} \equiv \left[\frac{k}{p} \right] \pmod{p}.$$

C 16. Suppose that p is an odd prime. Prove that

$$\sum_{j=0}^p \binom{p}{j} \binom{p+j}{j} \equiv 2^p + 1 \pmod{p^2}.$$

C 17. (Morley) Show that

$$(-1)^{\frac{p-1}{2}} \binom{p-1}{\frac{p-1}{2}} \equiv 4^{p-1} \pmod{p^3}$$

for all prime numbers p with $p \geq 5$.

C 18. Let n be a positive integer. Prove that n is prime if and only if

$$\binom{n-1}{k} \equiv (-1)^k \pmod{n}$$

for all $k \in \{0, 1, \dots, n-1\}$.

C 19. Prove that for $n \geq 2$,

$$\underbrace{2^{2^{\dots^2}}}_{n \text{ terms}} \equiv \underbrace{2^{2^{\dots^2}}}_{(n-1) \text{ terms}} \pmod{n}.$$

C 20. Show that, for any fixed integer $n \geq 1$, the sequence

$$2, 2^2, 2^{2^2}, 2^{2^{2^2}}, \dots \pmod{n}$$

is eventually constant.

C 21. Somebody incorrectly remembered Fermat's little theorem as saying that the congruence $a^{n+1} \equiv a \pmod{n}$ holds for all a if n is prime. Describe the set of integers n for which this property is in fact true.

C 22. Characterize the set of positive integers n such that, for all integers a , the sequence a, a^2, a^3, \dots is periodic modulo n .

C 23. Show that there exists a composite number n such that $a^n \equiv a \pmod{n}$ for all $a \in \mathbf{Z}$.

C 24. Let p be a prime number of the form $4k + 1$. Suppose that $2p + 1$ is prime. Show that there is no $k \in \mathbf{N}$ with $k < 2p$ and $2^k \equiv 1 \pmod{2p + 1}$.

C 25. During a break, n children at school sit in a circle around their teacher to play a game. The teacher walks clockwise close to the children and hands out candies to some of them according to the following rule. He selects one child and gives him a candy, then he skips the next child and gives a candy to the next one, then he skips 2 and gives a candy to the next one, then he skips 3, and so on. Determine the values of n for which eventually, perhaps after many rounds, all children will have at least one candy each.

C 26. Suppose that $m > 2$, and let P be the product of the positive integers less than m that are relatively prime to m . Show that $P \equiv -1 \pmod{m}$ if $m = 4, p^n$, or $2p^n$, where p is an odd prime, and $P \equiv 1 \pmod{m}$ otherwise.

C 27. Let Γ consist of all polynomials in x with integer coefficients. For f and g in Γ and m a positive integer, let $f \equiv g \pmod{m}$ mean that every coefficient of $f - g$ is an integral multiple of m . Let n and p be positive integers with p prime. Given that f, g, h, r and s are in Γ with $rf + sg \equiv 1 \pmod{p}$ and $fg \equiv h \pmod{p}$, prove that there exist F and G in Γ with $F \equiv f \pmod{p}$, $G \equiv g \pmod{p}$, and $FG \equiv h \pmod{p^n}$.

C 28. Determine the number of integers $n \geq 2$ for which the congruence

$$x^{2^5} \equiv x \pmod{n}$$

is true for all integers x .

C 29. Let n_1, \dots, n_k and a be positive integers which satisfy the following conditions :

- i) for any $i \neq j$, $(n_i, n_j) = 1$,
- ii) for any i , $a^{n_i} \equiv 1 \pmod{n_i}$, and
- iii) for any i , $n_i \nmid a - 1$.

Show that there exist at least $2^{k+1} - 2$ integers $x > 1$ with $a^x \equiv 1 \pmod{x}$.

C 30. Determine all positive integers $n \geq 2$ that satisfy the following condition ; For all integers a, b relatively prime to n ,

$$a \equiv b \pmod{n} \iff ab \equiv 1 \pmod{n}.$$

C 31. Determine all positive integers n such that $xy+1 \equiv 0 \pmod{n}$ implies that $x + y \equiv 0 \pmod{n}$.

C 32. Let p be a prime number. Determine the maximal degree of a polynomial $T(x)$ whose coefficients belong to $\{0, 1, \dots, p - 1\}$, whose degree is less than p , and which satisfies

$$T(n) = T(m) \pmod{p} \implies n = m \pmod{p}$$

for all integers n, m .

C 33. Let a_1, \dots, a_k and m_1, \dots, m_k be integers $2 \leq m_1$ and $2m_i \leq m_{i+1}$ for $1 \leq i \leq k - 1$. Show that there are infinitely many integers x which do not satisfy any of congruences

$$x \equiv a_1 \pmod{m_1}, x \equiv a_2 \pmod{m_2}, \dots, x \equiv a_k \pmod{m_k}.$$

C 34. Show that 1994 divides $10^{900} - 2^{1000}$.

C 35. Determine the last three digits of

$$2003^{2002^{2001}}.$$

C 36. Prove that $1980^{1981^{1982}} + 1982^{1981^{1980}}$ is divisible by 1981^{1981} .

C 37. Every odd prime is of the form $p = 4n + 1$.

- (a) Show that n is a quadratic residue \pmod{p} .
- (b) Calculate the value $n^n \pmod{p}$.

6. PRIMES AND COMPOSITE NUMBERS

Wherever there is number, there is beauty. Proclus Diadochus

6.1. Composite Numbers.

D 1. Prove that the number $512^3 + 675^3 + 720^3$ is composite.

D 2. Let a, b, c, d be integers with $a > b > c > d > 0$. Suppose that $ac + bd = (b + d + a - c)(b + d - a + c)$. Prove that $ab + cd$ is not prime.

D 3. Find the sum of all distinct positive divisors of the number 104060401.

D 4. Prove that 1280000401 is composite.

D 5. Prove that $\frac{5^{125}-1}{5^{25}-1}$ is a composite number.

D 6. Find the factor of $2^{33} - 2^{19} - 2^{17} - 1$ that lies between 1000 and 5000.

D 7. Show that there exists a positive integer k such that $k \cdot 2^n + 1$ is composite for all $n \in \mathbf{N}_0$.

D 8. Show that for all integer $k > 1$, there are infinitely many natural numbers n such that $k \cdot 2^{2^n} + 1$ is composite.

D 9. Four integers are marked on a circle. On each step we simultaneously replace each number by the difference between this number and next number on the circle in a given direction (that is, the numbers a, b, c, d are replaced by $a - b, b - c, c - d, d - a$). Is it possible after 1996 such steps to have numbers a, b, c , and d such that the numbers $|bc - ad|$, $|ac - bd|$, and $|ab - cd|$ are primes ?

D 10. Represent the number $989 \cdot 1001 \cdot 1007 + 320$ as the product of primes.

D 11. In 1772 Euler discovered the curious fact that $n^2 + n + 41$ is prime when n is any of $0, 1, 2, \dots, 39$. Show that there exist 40 consecutive integer values of n for which this polynomial is not prime.

6.2. Prime Numbers.

D 12. Show that there are infinitely many primes.

D 13. Find all natural numbers n for which every natural number whose decimal representation has $n - 1$ digits 1 and one digit 7 is prime.

D 14. Prove that there do not exist polynomials P and Q such that

$$\pi(x) = \frac{P(x)}{Q(x)}$$

for all $x \in \mathbf{N}$.

D 15. Show that there exist two consecutive squares such that there are at least 1000 primes between them.

D 16. Prove that for any prime p in the interval $(n, \frac{4n}{3}]$, p divides

$$\sum_{j=0}^n \binom{n}{j}^4$$

D 17. Let a, b , and n be positive integers with $\gcd(a, b) = 1$. Without using Dirichlet's theorem⁵, show that there are infinitely many $k \in \mathbf{N}$ such that $\gcd(ak + b, n) = 1$.

D 18. Without using Dirichlet's theorem, show that there are infinitely many primes ending in the digit 9.

D 19. Let p be an odd prime. Without using Dirichlet's theorem, show that there are infinitely many primes of the form $2pk + 1$.

D 20. Verify that, for each $r \geq 1$, there are infinitely many primes p with $p \equiv 1 \pmod{2^r}$.

D 21. Prove that if p is a prime, then $p^p - 1$ has a prime factor that is congruent to 1 modulo p .

D 22. Let p be a prime number. Prove that there exists a prime number q such that for every integer n , $n^p - p$ is not divisible by q .

D 23. Let $p_1 = 2, p_2 = 3, p_3 = 5, \dots, p_n$ be the first n prime numbers, where $n \geq 3$. Prove that

$$\frac{1}{p_1^2} + \frac{1}{p_2^2} + \dots + \frac{1}{p_n^2} + \frac{1}{p_1 p_2 \dots p_n} < \frac{1}{2}.$$

D 24. Let p_n be the n th prime : $p_1 = 2, p_2 = 3, p_3 = 5, \dots$. Show that the infinite series

$$\sum_{n=1}^{\infty} \frac{1}{p_n}$$

diverges.

D 25. Prove that $\log n \geq k \log 2$, where n is a natural number and k is the number of distinct primes that divide n .

D 26. Find the smallest prime which is not the difference (in some order) of a power of 2 and a power of 3.

D 27. Prove that for each positive integer n , there exist n consecutive positive integers none of which is an integral power of a prime number.

D 28. Show that $n^{\pi(2n) - \pi(n)} < 4^n$ for all positive integer n .

D 29. Let s_n denote the sum of the first n primes. Prove that for each n there exists an integer whose square lies between s_n and s_{n+1} .

⁵For any $a, b \in \mathbf{N}$ with $\gcd(a, b) = 1$, there are infinitely many primes of the form $ak + b$.

D 30. Given an odd integer $n > 3$, let k and t be the smallest positive integers such that both $kn + 1$ and tn are squares. Prove that n is prime if and only if both k and t are greater than $\frac{n}{4}$

D 31. Suppose n and r are nonnegative integers such that no number of the form $n^2 + r - k(k + 1)$ ($k \in \mathbf{N}$) equals to -1 or a positive composite number. Show that $4n^2 + 4r + 1$ is 1, 9 or prime.

D 32. Let $n \geq 5$ be an integer. Show that n is prime if and only if $n_i n_j \neq n_p n_q$ for every partition of n into 4 integers, $n = n_1 + n_2 + n_3 + n_4$, and for each permutation (i, j, p, q) of $(1, 2, 3, 4)$.

D 33. Prove that there are no positive integers a and b such that for all different primes p and q greater than 1000, the number $ap + bq$ is also prime.

D 34. Let p_n denote the n th prime number. For all $n \geq 6$, prove that

$$\pi(\sqrt{p_1 p_2 \cdots p_n}) > 2n.$$

D 35. There exists a block of 1000 consecutive positive integers containing no prime numbers, namely, $1001! + 2, 1001! + 3, \dots, 1001! + 1001$. Does there exist a block of 1000 consecutive positive integers containing exactly five prime numbers?

D 36. (S. Golomb) Prove that there are infinitely many twin primes if and only if there are infinitely many integers that cannot be written in any of the following forms :

$$6uv + u + v, \quad 6uv + u - v, \quad 6uv - u + v, \quad 6uv - u - v,$$

for some positive integers u and v .

D 37. It's known that there is always a prime between n and $2n - 7$ for all $n \geq 10$. Prove that, with the exception of 1, 4, and 6, every natural number can be written as the sum of distinct primes.

D 38. Prove that if $c > \frac{8}{3}$, then there exists a real numbers θ such that $[\theta^{c^n}]$ is prime for any positive integer n .

D 39. Let c be a nonzero real numbers. Suppose that

$$g(x) = c_0 x^r + c_1 x^{r-1} + \cdots + c_{r-1} x + c_r$$

is a polynomial with integer coefficients. Suppose that the roots of $g(x)$ are b_1, \dots, b_r . Let k be a given positive integer. Show that there is a prime p such that

$$p > k, |c|, |c_r|$$

and, moreover if t is a real between 0 and 1, and j is one of $1, \dots, r$, then

$$|(c^r b_j g(tb_j))^p e^{(1-t)b}| < \frac{(p-1)!}{2r}.$$

Furthermore, if

$$f(x) = \frac{e^{rp-1} x^{p-1} (g(x))^p}{(p-1)!}$$

then

$$\left| \sum_{j=1}^r \int_0^1 e^{(1-t)b_j} f(tb_j) dt \right| \leq \frac{1}{2}.$$

D 40. Prove that there do not exist eleven primes, all less than 20000, which can form an arithmetic progression.

D 41. (G. H. Hardy) Let n be a positive integer. Show that n is prime if and only if

$$\lim_{r \rightarrow \infty} \lim_{s \rightarrow \infty} \lim_{t \rightarrow \infty} \sum_{u=0}^s \left(1 - \left(\cos \frac{(u!)^r \pi}{n} \right)^2 t \right) = n.$$

7. RATIONAL AND IRRATIONAL NUMBERS

God made the integers, all else is the work of man. Leopold Kronecker

7.1. Rational Numbers.

E 1. Suppose that a rectangle with sides a and b is arbitrarily cut into squares with sides x_1, \dots, x_n . Show that $\frac{x_i}{a} \in \mathbb{Q}$ and $\frac{x_i}{b} \in \mathbb{Q}$ for all $i \in \{1, \dots, n\}$.

E 2. Find all x and y which are rational multiples of π with $0 < x < y < \frac{\pi}{2}$ and $\tan x + \tan y = 2$.

E 3. Let α be a rational number with $0 < \alpha < 1$ and $\cos(3\pi\alpha) + 2\cos(2\pi\alpha) = 0$. Prove that $\alpha = \frac{2}{3}$.

E 4. Suppose that $\tan \alpha = \frac{p}{q}$, where p and q are integers and $q \neq 0$. Prove the number $\tan \beta$ for which $\tan 2\beta = \tan 3\alpha$ is rational only when $p^2 + q^2$ is the square of an integer.

E 5. Prove that there is no positive rational number x such that

$$x^{[x]} = \frac{9}{2}.$$

E 6. Let x, y, z non-zero real numbers such that xy, yz, zx are rational.

(a) Show that the number $x^2 + y^2 + z^2$ is rational.

(b) If the number $x^3 + y^3 + z^3$ is also rational, show that x, y, z are rational.

E 7. If x is a positive rational number, show that x can be uniquely expressed in the form

$$x = a_1 + \frac{a_2}{2!} + \frac{a_3}{3!} + \dots,$$

where a_1, a_2, \dots are integers, $0 \leq a_n \leq n - 1$, for $n > 1$, and the series terminates. Show also that x can be expressed as the sum of reciprocals of different integers, each of which is greater than 10^6 .

E 8. Find all polynomials W with real coefficients possessing the following property : if $x + y$ is a rational number, then $W(x) + W(y)$ is rational.

E 9. Prove that every positive rational number can be represented in the form

$$\frac{a^3 + b^3}{c^3 + d^3}$$

for some positive integers a, b, c , and d .

E 10. The set S is a finite subset of $[0, 1]$ with the following property : for all $s \in S$, there exist $a, b \in S \cup \{0, 1\}$ with $a, b \neq s$ such that $s = \frac{a+b}{2}$. Prove that all the numbers in S are rational.

E 11. Let $S = \{x_0, x_1, \dots, x_n\} \subset [0, 1]$ be a finite set of real numbers with $x_0 = 0$ and $x_1 = 1$, such that every distance between pairs of elements occurs at least twice, except for the distance 1. Prove that all of the x_i are rational.

E 12. Does there exist a circle and an infinite set of points on it such that the distance between any two points of the set is rational ?

E 13. Prove that numbers of the form

$$\frac{a_1}{1!} + \frac{a_2}{2!} + \frac{a_3}{3!} + \dots,$$

where $0 \leq a_i \leq i - 1$ ($i = 2, 3, 4, \dots$) are rational if and only if starting from some i on all the a_i 's are either equal to 0 (in which case the sum is finite) or all are equal to $i - 1$.

E 14. Let k and m be positive integers. Show that

$$S(m, k) = \sum_{n=1}^{\infty} \frac{1}{n(mn + k)}$$

is rational if and only if m divides k .

E 15. Find all rational numbers k such that $0 \leq k \leq \frac{1}{2}$ and $\cos k\pi$ is rational.

E 16. Prove that for any distinct rational numbers of a, b, c , the number

$$\frac{1}{(b - c)^2} + \frac{1}{(c - a)^2} + \frac{1}{(a - b)^2}$$

is the square of some rational number.

7.2. Irrational Numbers.

E 17. Find the smallest positive integer n such that

$$0 < n^{\frac{1}{4}} - [n^{\frac{1}{4}}] < 0.00001.$$

E 18. Prove that for any positive integers a and b

$$\left| a\sqrt{2} - b \right| > \frac{1}{2(a + b)}.$$

E 19. Prove that there exist positive integers m and n such that

$$\left| \frac{m^2}{n^3} - \sqrt{2001} \right| < \frac{1}{10^8}.$$

E 20. Let a, b, c be integers, not all zero and each of absolute value less than one million. Prove that

$$\left| a + b\sqrt{2} + c\sqrt{3} \right| > \frac{1}{10^{21}}.$$

E 21. Let a, b, c be integers, not all equal to 0. Show that

$$\frac{1}{4a^2 + 3b^2 + 2c^2} \leq |\sqrt[3]{4a} + \sqrt[3]{2b} + c|.$$

E 22. (Hurwitz) Prove that for any irrational number ξ , there are infinitely many rational numbers $\frac{m}{n}$ $((m, n) \in \mathbf{Z} \times \mathbf{N})$ such that

$$\left| \xi - \frac{n}{m} \right| < \frac{1}{\sqrt{5}m^2}.$$

E 23. Show that π is irrational.

E 24. Show that $e = \sum_{n=0}^{\infty} \frac{1}{n!}$ is irrational.

E 25. Show that $\cos \frac{\pi}{7}$ is irrational.

E 26. Show that $\frac{1}{\pi} \arccos\left(\frac{1}{\sqrt{2003}}\right)$ is irrational.

E 27. Show that $\cos 1^\circ$ is irrational.

E 28. An integer-sided triangle has angles $p\theta$ and $q\theta$, where p and q are relatively prime integers. Prove that $\cos \theta$ is irrational.

E 29. It is possible to show that $\csc \frac{3\pi}{29} - \csc \frac{10\pi}{29} = 1.999989433\dots$. Prove that there are no integers j, k, n with odd n satisfying $\csc \frac{j\pi}{n} - \csc \frac{k\pi}{n} = 2$.

E 30. For which angles θ , a rational number of degrees, is it the case that $\tan^2\theta + \tan^2 2\theta$ is irrational?

E 31. (K. Mahler, 1953) Prove that for any $p, q \in \mathbf{N}$ with $q > 1$ the following inequality holds.⁶

$$\left| \pi - \frac{p}{q} \right| \geq q^{-42}$$

E 32. For each integer $n \geq 1$, prove that there is a polynomial $P_n(x)$ with rational coefficients such that

$$x^{4n}(1-x)^{4n} = (1+x)^2 P_n(x) + (-1)^n 4^n.$$

Define the rational number a_n by

$$a_n = \frac{(-1)^{n-1}}{4^{n-1}} \int_0^1 P_n(x) dx, \quad n = 1, 2, \dots$$

Prove that a_n satisfies the inequality

$$|\pi - a_n| < \frac{1}{4^{5n-1}}, \quad n = 1, 2, \dots$$

E 33. (K. Alladi, M. Robinson, 1979) Suppose that $p, q \in \mathbf{N}$ satisfy the inequality $e(\sqrt{p+q} - \sqrt{q})^2 < 1$.⁷ Show that $\ln\left(1 + \frac{p}{q}\right)$ is irrational.

⁶This is a deep theorem in *transcendental number theory*. Note that it follows from this result that π is irrational! In fact, it's known that for sufficiently large q , the exponent 42 can be replaced by 30. Here is a similar result due to A. Baker: For any rationals $\frac{p}{q}$, one has $|\ln 2 - \frac{p}{q}| \geq 10^{-100000} q^{-12.5}$. [AI, pp. 106]

⁷Here, $e = \sum_{n \geq 0} \frac{1}{n!}$.

E 34. Show that the cube roots of three distinct primes cannot be terms in an arithmetic progression.

E 35. Let n be an integer greater than or equal to 3. Prove that there is a set of n points in the plane such that the distance between any two points is irrational and each set of three points determines a non-degenerate triangle with a rational area.

E 36. You are given three lists A , B , and C . List A contains the numbers of the form 10^k in base 10, with k any integer greater than or equal to 1. Lists B and C contain the same numbers translated into base 2 and 5 respectively:

A	B	C
10	1010	20
100	1100100	400
1000	1111101000	13000
\vdots	\vdots	\vdots

Prove that for every integer $n > 1$, there is exactly one number in exactly one of the lists B or C that has exactly n digits.

E 37. (Beatty) Prove that if α and β are positive irrational numbers satisfying $\frac{1}{\alpha} + \frac{1}{\beta} = 1$, then the sequences

$$[\alpha], [2\alpha], [3\alpha], \dots$$

and

$$[\beta], [2\beta], [3\beta], \dots$$

together include every positive integer exactly once.

E 38. For a positive real number α , define

$$S(\alpha) = \{[n\alpha] \mid n = 1, 2, 3, \dots\}.$$

Prove that \mathbf{N} cannot be expressed as the disjoint union of three sets $S(\alpha)$, $S(\beta)$, and $S(\gamma)$.

E 39. Let $f(x) = \prod_{n=1}^{\infty} (1 + \frac{x}{2^n})$. Show that at the point $x = 1$, $f(x)$ and all its derivatives are irrational.

E 40. Let $\{a_n\}_{n \geq 1}$ be a sequence of positive numbers such that

$$a_{n+1}^2 = a_n + 1, \quad n \in \mathbf{N}.$$

Show that the sequence contains an irrational number.

E 41. Show that $\tan(\frac{\pi}{m})$ is irrational for all positive integers $m \geq 5$.

E 42. Prove that if $g \geq 2$ is an integer, then two series

$$\sum_{n=0}^{\infty} \frac{1}{g^{n^2}} \quad \text{and} \quad \sum_{n=0}^{\infty} \frac{1}{g^{n!}}$$

both converge to irrational numbers.

E 43. Let $1 < a_1 < a_2 < \dots$ be a sequence of positive integers. Show that

$$\frac{2^{a_1}}{a_1!} + \frac{2^{a_2}}{a_2!} + \frac{2^{a_3}}{a_3!} + \dots$$

is irrational.

E 44. (N. Agahanov) Do there exist real numbers a and b such that

(1) $a + b$ is rational and $a^n + b^n$ is irrational for all $n \in \mathbf{N}$ with $n \geq 2$?

(2) $a + b$ is irrational and $a^n + b^n$ is rational for all $n \in \mathbf{N}$ with $n \geq 2$?

E 45. Let $p(x) = x^3 + a_1x^2 + a_2x + a_3$ have rational coefficients and have roots $r_1, r_2,$ and r_3 . If $r_1 - r_2$ is rational, must $r_1, r_2,$ and r_3 be rational ?

E 46. Let $\alpha = 0.d_1d_2d_3\dots$ be a decimal representation of a real number between 0 and 1. Let r be a real number with $|r| < 1$.

(a) If α and r are rational, must $\sum_{i=1}^{\infty} d_i r^i$ be rational ?

(b) If $\sum_{i=1}^{\infty} d_i r^i$ and r are rational, α must be rational ?

8. DIOPHANTINE EQUATIONS I

In the margin of his copy of Diophantus' *Arithmetica*, Pierre de Fermat wrote : *To divide a cube into two other cubes, a fourth power or in general any power whatever into two powers of the same denomination above the second is impossible, and I have assuredly found an admirable proof of this, but the margin is too narrow to contain it.*

F 1. *One of Euler's conjecture⁸ was disproved in the 1980s by three American Mathematicians⁹ when they showed that there is a positive integer n such that*

$$n^5 = 133^5 + 110^5 + 84^5 + 27^5.$$

Find the value of n .

F 2. *The number 21982145917308330487013369 is the thirteenth power of a positive integer. Which positive integer?*

F 3. *Does there exist a solution to the equation*

$$x^2 + y^2 + z^2 + u^2 + v^2 = xyzuv - 65$$

in integers x, y, z, u, v greater than 1998?

F 4. *Find all pairs (x, y) of positive rational numbers such that $x^2 + 3y^2 = 1$.*

F 5. *Find all pairs (x, y) of rational numbers such that $y^2 = x^3 - 3x + 2$.*

F 6. *Show that there are infinitely many pairs (x, y) of rational numbers such that $x^3 + y^3 = 9$.*

F 7. *Determine all pairs (x, y) of positive integers satisfying the equation*

$$(x + y)^2 - 2(xy)^2 = 1.$$

F 8. *Show that the equation*

$$x^3 + y^3 + z^3 + t^3 = 1999$$

has infinitely many integral solutions.¹⁰

F 9. *Determine all integers a for which the equation*

$$x^2 + axy + y^2 = 1$$

has infinitely many distinct integer solutions x, y .

⁸In 1769, Euler, by generalizing Fermat's Last Theorem, conjectured that "it is impossible to exhibit three fourth powers whose sum is a fourth power", "four fifth powers whose sum is a fifth power, and similarly for higher powers" [Rs]

⁹L. J. Lander, T. R. Parkin, and J. L. Selfridge

¹⁰More generally, the following result is known : let n be an integer, then the equation $x^3 + y^3 + z^3 + w^3 = n$ has infinitely many integral solutions (x, y, z, w) if there can be found one solution $(x, y, z, w) = (a, b, c, d)$ with $(a + b)(c + d)$ negative and with either $a \neq b$ and $c \neq d$. [Eb2, pp.90]

F 10. Prove that there are unique positive integers a and n such that

$$a^{n+1} - (a+1)^n = 2001.$$

F 11. Find all $(x, y, n) \in \mathbf{N}^3$ such that $\gcd(x, n+1) = 1$ and $x^n + 1 = y^{n+1}$.

F 12. Find all $(x, y, z) \in \mathbf{N}^3$ such that $x^4 - y^4 = z^2$.

F 13. Find all pairs (x, y) of positive integers that satisfy the equation ¹¹

$$y^2 = x^3 + 16.$$

F 14. Show that the equation $x^2 + y^5 = z^3$ has infinitely many solutions in integers x, y, z for which $xyz \neq 0$.

F 15. Prove that there are no integers x and y satisfying $x^2 = y^5 - 4$.

F 16. Find all pairs (a, b) of different positive integers that satisfy the equation $W(a) = W(b)$, where $W(x) = x^4 - 3x^3 + 5x^2 - 9x$.

F 17. Find all positive integers n for which the equation

$$a + b + c + d = n\sqrt{abcd}$$

has a solution in positive integers.

F 18. Determine all positive integer solutions (x, y, z, t) of the equation

$$(x+y)(y+z)(z+x) = xyzt$$

for which $\gcd(x, y) = \gcd(y, z) = \gcd(z, x) = 1$.

F 19. Find all $(x, y, z, n) \in \mathbf{N}^4$ such that $x^3 + y^3 + z^3 = nx^2y^2z^2$.

F 20. Determine all positive integers n for which the equation

$$x^n + (2+x)^n + (2-x)^n = 0$$

has an integer as a solution.

F 21. Prove that the equation

$$6(6a^2 + 3b^2 + c^2) = 5n^2$$

has no solutions in integers except $a = b = c = n = 0$.

F 22. Find all integers a, b, c, x, y, z such that

$$a + b + c = xyz, \quad x + y + z = abc, \quad a \geq b \geq c \geq 1, \quad x \geq y \geq z \geq 1.$$

F 23. Find all $(x, y, z) \in \mathbf{Z}^3$ such that $x^3 + y^3 + z^3 = x + y + z = 3$.

¹¹It's known that there are (infinitely) many integers k so that the equation $y^2 = x^3 + k$ has no integral solutions. For example, if k has the form $k = (4n-1)^3 - 4m^2$, where m and n are integers such that no prime $p \equiv -1 \pmod{4}$ divides m , then the equation $y^2 = x^3 + k$ has no integral solutions. For a proof, see [Tma, pp. 191].

F 24. Prove that if n is a positive integer such that the equation

$$x^3 - 3xy^2 + y^3 = n.$$

has a solution in integers (x, y) , then it has at least three such solutions.

Show that the equation has no solutions in integers when $n = 2891$.

F 25. What is the smallest positive integer t such that there exist integers x_1, x_2, \dots, x_t with

$$x_1^3 + x_2^3 + \dots + x_t^3 = 2002^{2002} \quad ?$$

F 26. Solve in integers the following equation

$$n^{2002} = m(m+n)(m+2n) \cdots (m+2001n).$$

F 27. Prove that there exist infinitely many positive integers n such that $p = nr$, where p and r are respectively the semi-perimeter and the inradius of a triangle with integer side lengths.

F 28. Let a, b, c be positive integers such that a and b are relatively prime and c is relatively prime either to a and b . Prove that there exist infinitely many triples (x, y, z) of distinct positive integers such that

$$x^a + y^b = z^c.$$

F 29. Find all pairs of integers (x, y) satisfying the equality

$$y(x^2 + 36) + x(y^2 - 36) + y^2(y - 12) = 0$$

F 30. Let a, b, c be given integers $a > 0$, $ac - b^2 = P = P_1 P_2 \cdots P_n$, where P_1, \dots, P_n are (distinct) prime numbers. Let $M(n)$ denote the number of pairs of integers (x, y) for which $ax^2 + bxy + cy^2 = n$. Prove that $M(n)$ is finite and $M(n) = M(p^k \cdot n)$ for every integers $k \geq 0$.

F 31. Determine integer solutions of the system

$$2uv - xy = 16,$$

$$xv - yu = 12.$$

F 32. Let n be a natural number. Solve in whole numbers the equation

$$x^n + y^n = (x - y)^{n+1}.$$

F 33. Does there exist an integer such that its cube is equal to $3n^2 + 3n + 7$, where n is integer?

F 34. Are there integers m and n such that $5m^2 - 6mn + 7n^2 = 1985$?

F 35. Find all cubic polynomials $x^3 + ax^2 + bx + c$ admitting the rational numbers a, b and c as roots.

F 36. Prove that the equation $a^2 + b^2 = c^2 + 3$ has infinitely many integer solutions (a, b, c) .

F 37. Prove that for each positive integer n there exist odd positive integers x_n and y_n such that $x_n^2 + 7y_n^2 = 2^n$.

F 38. Suppose that p is an odd prime such that $2p + 1$ is also prime. Show that the equation $x^p + 2y^p + 5z^p = 0$ has no solutions in integers.

F 39. Let A, B, C, D, E be integers $B \neq 0$ and $F = AD^2 - BCD + B^2E \neq 0$. Prove that the number N of pairs of integers (x, y) such that

$$Ax^2 + Bxy + Cx + Dy + E = 0,$$

satisfies $N \leq 2d(|F|)$, where $d(n)$ denotes the number of positive divisors of positive integer n .

F 40. Determine all pairs of rational numbers (x, y) such that

$$x^3 + y^3 = x^2 + y^2.$$

F 41. Suppose that $A = 1, 2$, or 3 . Let a and b be relatively prime integers such that $a^2 + Ab^2 = s^3$ for some integer s . Then, there are integers u and v such that $s = u^2 + Av^2$, $a = u^3 - 3Avu^2$, and $b = 3u^2v - Av^3$.

F 42. Find all integers a for which $x^3 - x + a$ has three integer roots.

F 43. Find all solutions in integers of $x^3 + 2y^3 = 4z^3$.

F 44. For a $n \in \mathbf{N}$, show that the number of integral solutions (x, y) of

$$x^2 + xy + y^2 = n$$

is finite and a multiple of 6.

F 45. (Fermat) Show that there cannot be four squares in arithmetical progression.

F 46. (Gauss) Let a, b, c, d, e, f be integers such that $b^2 - 4ac > 0$ is not a perfect square and $4acf + bde - ae^2 - cd^2 - fb^2 \neq 0$. Let

$$f(x, y) = ax^2 + bxy + cy^2 + dx + ey + f$$

Suppose that $f(x, y) = 0$ has an integral solution. Show that $f(x, y) = 0$ has infinitely many integral solutions.

F 47. Show that the equation $x^4 + y^4 + 4z^4 = 1$ has infinitely many rational solutions.

F 48. Solve the equation $x^2 + 7 = 2^n$ in integers.

F 49. Show that the only solutions of the equation $x^3 - 3xy^2 - y^3 = 1$ are given by $(x, y) = (1, 0), (0, -1), (-1, 1), (1, -3), (-3, 2), (2, 1)$.

F 50. Show that the equation $y^2 = x^3 + 2a^3 - 3b^2$ has no solution in integers if $ab \neq 0$, $a \not\equiv 1 \pmod{3}$, $3 \nmid b$, a is odd if b is even, and $p = t^2 + 27u^2$ is soluble in integers t and u of $p \mid a$ and $p \equiv 1 \pmod{3}$.

F 51. Prove that the product of five consecutive integers is never a perfect square.

F 52. Do there exist two right-angled triangles with integer length sides that have the lengths of exactly two sides in common?

F 53. Suppose that a, b , and p are integers such that $b \equiv 1 \pmod{4}$, $p \equiv 3 \pmod{4}$, p is prime, and if q is any prime divisor of a such that $q \equiv 3 \pmod{4}$, then $q^p | a^2$ and $p \nmid q - 1$ (if $q = p$, then also $q | b$). Show that the equation

$$x^2 + 4a^2 = y^p - b^p$$

has no solutions in integers.

F 54. Show that the number of integral-sided right triangles whose ratio of area to semi-perimeter is p^m , where p is a prime and m is an integer, is $m + 1$ if $p = 2$ and $2m + 1$ if $p \neq 2$.

9. DIOPHANTINE EQUATIONS II

The positive integers stand there, a continual and inevitable challenge to the curiosity of every healthy mind. Godfrey Harold Hardy

G 1. *Given that*

$$34! = 95232799cd96041408476186096435ab000000_{(10)},$$

determine the digits a, b, c , and d .

G 2. *Prove that the equation $(x_1 - x_2)(x_2 - x_3)(x_3 - x_4)(x_4 - x_5)(x_5 - x_6)(x_6 - x_7)(x_7 - x_1) = (x_1 - x_3)(x_2 - x_4)(x_3 - x_5)(x_4 - x_6)(x_5 - x_7)(x_6 - x_1)(x_7 - x_2)$ has a solution in natural numbers where all x_i are different.*

G 3. *(P. Erdős) Show that the equation $\binom{n}{k} = m^l$ has no integral solution with $l \geq 2$ and $4 \leq k \leq n - 4$.*

G 4. *Solve in positive integers the equation $10^a + 2^b - 3^c = 1997$.*

G 5. *Solve the equation $28^x = 19^y + 87^z$, where x, y, z are integers.*

G 6. *Show that the equation $x^7 + y^7 = 1998^z$ has no solution in positive integers.*

G 7. *Solve the equation $2^x - 5 = 11^y$ in positive integers.*

G 8. *Solve the equation $7^x - 3^y = 4$ in positive integers.*

G 9. *Show that $|12^m - 5^n| \geq 7$ for all $m, n \in \mathbf{N}$.*

G 10. *Show that there is no positive integer k for which the equation*

$$(n - 1)! + 1 = n^k$$

is true when n is greater than 5.

G 11. *Determine all pairs (x, y) of integers such that*

$$(19a + b)^{18} + (a + b)^{18} + (19b + a)^{18}$$

is a positive square.

G 12. *Let b be a positive integer. Determine all 200-tuple integers of non-negative integers $(a_1, a_2, \dots, a_{200})$ satisfying*

$$\sum_{j=1}^n a_j^{a_j} = 2002b^b.$$

G 13. *Is there a positive integers m such that the equation*

$$\frac{1}{a} + \frac{1}{b} + \frac{1}{c} + \frac{1}{abc} = \frac{m}{a + b + c}$$

has infinitely many solutions in positive integers a, b, c ?

G 14. Consider the system

$$\begin{aligned}x + y &= z + u, \\ 2xy &= zu.\end{aligned}$$

Find the greatest value of the real constant m such that $m \leq \frac{x}{y}$ for any positive integer solution (x, y, z, u) of the system, with $x \geq y$.

G 15. Determine all positive rational numbers $r \neq 1$ such that $r^{\frac{1}{r-1}}$ is rational.

G 16. Show that the equation $\{x^3\} + \{y^3\} = \{z^3\}$ has infinitely many rational non-integer solutions.

G 17. Let n be a positive integer. Prove that the equation

$$x + y + \frac{1}{x} + \frac{1}{y} = 3n$$

does not have solutions in positive rational numbers.

G 18. Find all pairs (x, y) of positive rational numbers such that $x^y = y^x$

G 19. Find all pairs (a, b) of positive integers that satisfy the equation

$$a^{b^2} = b^a.$$

G 20. Find all pairs (a, b) of positive integers that satisfy the equation

$$a^{a^a} = b^b.$$

G 21. Let a, b , and x be positive integers such that $x^{a+b} = a^{bb}$. Prove that $a = x$ and $b = x^x$.

G 22. Find all pairs (m, n) of integers that satisfy the equation

$$(m - n)^2 = \frac{4mn}{m + n - 1}.$$

G 23. Find all pairwise relatively prime positive integers l, m, n such that

$$(l + m + n) \left(\frac{1}{l} + \frac{1}{m} + \frac{1}{n} \right)$$

is an integer.

G 24. Let x, y , and z be integers with $z > 1$. Show that

$$(x + 1)^2 + (x + 2)^2 + \cdots + (x + 99)^2 \neq y^z.$$

G 25. Find all values of the positive integers m and n for which

$$1! + 2! + 3! + \cdots + n! = m^2$$

G 26. Prove that if a, b, c, d are integers such that $d = (a + 2^{\frac{1}{3}}b + 2^{\frac{2}{3}}c)^2$ then d is a perfect square (i. e. is the square of an integer).

G 27. Find a pair of relatively prime four digit natural numbers A and B such that for all natural numbers m and n , $|A^m - B^n| \geq 400$.

G 28. Find all triples (a, b, c) of positive integers to the equation

$$a!b! = a! + b! + c!.$$

G 29. Find all pairs (a, b) of positive integers such that

$$(\sqrt[3]{a} + \sqrt[3]{b} - 1)^2 = 49 + 20\sqrt[3]{6}.$$

G 30. For what positive numbers a is

$$\sqrt[3]{2 + \sqrt{a}} + \sqrt[3]{2 - \sqrt{a}}$$

an integer ?

G 31. Find all integer solutions to $2(x^5 + y^5 + 1) = 5xy(x^2 + y^2 + 1)$.

G 32. A triangle with integer sides is called Heronian if its area is an integer. Does there exist a Heronian triangle whose sides are the arithmetic, geometric and harmonic means of two positive integers ?

G 33. What is the smallest perfect square that ends in 9009?

G 34. (Leo Moser) Show that the Diophantine equation

$$\frac{1}{x_1} + \frac{1}{x_2} + \cdots + \frac{1}{x_n} + \frac{1}{x_1 x_2 \cdots x_n} = 1$$

has at least one solution for every positive integers n .

G 35. Prove that the number $99999 + 111111\sqrt{3}$ cannot be written in the form $(A + B\sqrt{3})^2$, where A and B are integers.

G 36. Find all triples of positive integers (x, y, z) such that

$$(x + y)(1 + xy) = 2^z.$$

G 37. If R and S are two rectangles with integer sides such that the perimeter of R equals the area of S and the perimeter of S equals the area of R , call R and S are amicable pair of rectangles. Find all amicable pairs of rectangles.

10. FUNCTIONS IN NUMBER THEORY

Gauss once said "Mathematics is the queen of the sciences and number theory is the queen of mathematics." If this be true we may add that the Disquisitiones is the Magna Charta of number theory. M. Cantor

10.1. Floor Function and Fractional Part Function.

H 1. Let α be the positive root of the equation $x^2 = 1991x + 1$. For natural numbers m and n define

$$m * n = mn + [\alpha m][\alpha n],$$

where $[x]$ is the greatest integer not exceeding x . Prove that for all natural numbers p, q , and r ,

$$(p * q) * r = p * (q * r).$$

H 2. Prove that for any positive integer n ,

$$\left[\frac{n}{3}\right] + \left[\frac{n+2}{6}\right] + \left[\frac{n+4}{6}\right] = \left[\frac{n}{2}\right] + \left[\frac{n+3}{6}\right]$$

H 3. Prove that for any positive integer n ,

$$\left[\frac{n+1}{2}\right] + \left[\frac{n+2}{4}\right] + \left[\frac{n+4}{8}\right] + \left[\frac{n+8}{16}\right] + \dots = n$$

H 4. Show that for all positive integers n ,

$$[\sqrt{n} + \sqrt{n+1}] = [\sqrt{4n+1}] = [\sqrt{4n+2}] = [\sqrt{4n+3}].$$

H 5. Find all real numbers α for which the equality

$$[\sqrt{n} + \sqrt{n+\alpha}] = [\sqrt{4n+1}]$$

holds for all positive integers n .

H 6. Prove that for all positive integers n ,

$$[\sqrt{n} + \sqrt{n+1} + \sqrt{n+2}] = [\sqrt{9n+8}].$$

H 7. Prove that for all positive integers n ,

$$[n^{\frac{1}{3}} + (n+1)^{\frac{1}{3}}] = [(8n+3)^{\frac{1}{3}}]$$

H 8. Prove that $[n^{\frac{1}{3}} + (n+1)^{\frac{1}{3}} + (n+2)^{\frac{1}{3}}] = [(27n+26)^{\frac{1}{3}}]$ for all positive integer n .

H 9. Show that for all positive integers m and n ,

$$\gcd(m, n) = m + n - mn + 2 \sum_{k=0}^{m-1} \left[\frac{kn}{m} \right].$$

H 10. Show that for all primes p ,

$$\sum_{k=1}^{p-1} \left[\frac{k^3}{p} \right] = \frac{(p+1)(p-1)(p-2)}{4}.$$

H 11. Let p be a prime number of the form $4k+1$. Show that

$$\sum_{i=1}^{p-1} \left(\left[\frac{2i^2}{p} \right] - 2 \left[\frac{i^2}{p} \right] \right) = \frac{p-1}{2}.$$

H 12. Let $p = 4k+1$ be a prime. Show that

$$\sum_{i=1}^k \left[\sqrt{ip} \right] = \frac{p^2-1}{12}.$$

H 13. Suppose that $n \geq 2$. Prove that

$$\sum_{k=2}^n \left[\frac{n^2}{k} \right] = \sum_{k=n+1}^{n^2} \left[\frac{n^2}{k} \right]$$

H 14. Let a, b, n be positive integers with $\gcd(a, b) = 1$. Prove that

$$\sum_k \left\{ \frac{ak+b}{n} \right\} = \frac{n-1}{2},$$

where k runs through a complete system of residues modulo m .

H 15. Find the total number of different integer values the function

$$f(x) = [x] + [2x] + \left[\frac{5x}{3} \right] + [3x] + [4x]$$

takes for real numbers x with $0 \leq x \leq 100$.

H 16. Prove or disprove that there exists a positive real number u such that $[u^n] - n$ is an even integer for all positive integer n .

H 17. Determine all real numbers a such that

$$4[an] = n + [a[an]] \text{ for all } n \in \mathbf{N}$$

H 18. Do there exist irrational numbers $a, b > 1$ and $[a^m]$ differs $[b^n]$ for any two positive integers m and n ?

H 19. Let a, b, c , and d be real numbers. Suppose that $[na] + [nb] = [nc] + [nd]$ for all positive integers n . Show that at least one of $a+b$, $a-c$, $a-d$ is an integer.

H 20. (S. Reznichenko) Find all integer solutions of the equation

$$\left[\frac{x}{1!} \right] + \left[\frac{x}{2!} \right] + \cdots + \left[\frac{x}{10!} \right] = 1001.$$

10.2. Euler phi Function.

H 21. Let n be an integer with $n \geq 2$. Show that $\phi(2^n - 1)$ is divisible by n .

H 22. (Gauss) Show that for all $n \in \mathbf{N}$,

$$n = \sum_{d|n} \phi(d).$$

H 23. If p is a prime and n an integer such that $1 < n \leq p$, then

$$\phi\left(\sum_{k=0}^{p-1} n^k\right) \equiv 0 \pmod{p}.$$

H 24. Let m, n be positive integers. Prove that, for some positive integer a , each of $\phi(a), \phi(a+1), \dots, \phi(a+n)$ is a multiple of m .

H 25. If n is composite, prove that $\phi(n) \leq n - \sqrt{n}$.

H 26. Show that if m and n are relatively prime positive integers, then $\phi(5^m - 1) \neq 5^n - 1$.

H 27. Show that if the equation $\phi(x) = n$ has one solution it always has a second solution, n being given and x being the unknown.

H 28. Prove that for any δ greater than 1 and any positive number ϵ , there is an n such that $\left|\frac{\phi(n)}{n} - \delta\right| < \epsilon$.

H 29. (Schinzel, Sierpiński) Show that the set of all numbers $\frac{\phi(n+1)}{\phi(n)}$ is dense in the set of all positive reals.

H 30. (a) Show that if $n > 49$, then there are $a > 1$ and $b > 1$ such that $a + b = n$ and $\frac{\phi(a)}{a} + \frac{\phi(b)}{b} < 1$. (b) Show that if $n > 4$, then there are $a > 1$ and $b > 1$ such that $a + b = n$ and $\frac{\phi(a)}{a} + \frac{\phi(b)}{b} > 1$.

10.3. Divisor Functions.

H 31. Prove that $d(n^2 + 1)^2$ does not become monotonic from any given point onwards.

H 32. Determine all positive integers n such that $n = d(n)^2$.

H 33. Determine all positive integers k such that

$$\frac{d(n^2)}{d(n)} = k$$

for some $n \in \mathbf{N}$.

H 34. Find all positive integers n such that $d(n)^3 = 4n$.

H 35. Determine all positive integers for which $d(n) = \frac{n}{3}$ holds.

H 36. We say that an integer $m \geq 1$ is super-abundant if

$$\frac{\sigma(m)}{m} > \frac{\sigma(k)}{k},$$

for all $k \in \{1, 2, \dots, m-1\}$. Prove that there exists an infinite number of super-abundant numbers.

H 37. Let $\sigma(n)$ denote the sum of the positive divisors of the positive integer n . and $\phi(n)$ the Euler phi-function. Show that $\phi(n) + \sigma(n) \geq 2n$ for all positive integers n .

H 38. Prove that for any δ greater than 1 and any positive number ϵ , there is an n such that $\left| \frac{\sigma(n)}{n} - \delta \right| < \epsilon$.

H 39. Prove that $\sigma(n)\phi(n) < n^2$, but that there is a positive constant c such that $\sigma(n)\phi(n) \geq cn^2$ holds for all positive integers n .

H 40. Show that $\sigma(n) - d(m)$ is even for all positive integers m and n where m is the largest odd divisor of n .

H 41. Verify the Ramanujan sum

$$\sum_{d|\gcd(m,n)} d\mu\left(\frac{n}{d}\right) = \frac{\left(\frac{n}{\gcd(m,n)}\right)\phi(n)}{\phi\left(\frac{n}{\gcd(m,n)}\right)}.$$

H 42. Show that for any positive integer n ,

$$\frac{\sigma(n!)}{n!} \geq \sum_{k=1}^n \frac{1}{k}.$$

10.4. More Functions.

H 43. Ramanujan's tau Function¹² $\tau : \mathbf{N} \rightarrow \mathbf{Z}$ has the generating function

$$\sum_{n=1}^{\infty} \tau(n)x^n = x \prod_{n=1}^{\infty} (1-x^n)^{24},$$

i.e. the coefficients of x^n on the right hand side define $\tau(n)$.¹³

(1) Show that $\tau(mn) = \tau(m)\tau(n)$ for all $m, n \in \mathbf{N}$ with $\gcd(m, n) = 1$.¹⁴

(2) Show that $\tau(n) \equiv \sum_{d|n} d^{11} \pmod{691}$ for all $n \in \mathbf{N}$.¹⁵

H 44. For every natural number n , $Q(n)$ denote the sum of the digits in the decimal representation of n . Prove that there are infinitely many natural numbers k with $Q(3^k) > Q(3^{k+1})$.

H 45. Let $S(n)$ be the sum of all different natural divisors of an odd natural number $n > 1$ (including 1 and n). Prove that $S(n)^3 < n^4$.

¹²In 1947, Lehmer conjectured that $\tau(n) \neq 0$ for all $n \in \mathbf{N}$.

¹³ $\{\tau(n)|n \geq 1\} = \{1, -24, 252, -1472, \dots\}$. For more terms, see the first page!

¹⁴This Ramanujan's conjecture was proved by Mordell.

¹⁵This Ramanujan's conjecture was proved by Watson.

H 46. Let $((x)) = x - [x] - \frac{1}{2}$ if x is not an integer, and let $((x)) = 0$ otherwise. If n and k are integers, with $n > 0$, prove that

$$\left(\left(\frac{k}{n}\right)\right) = -\frac{1}{2n} \sum_{m=1}^{n-1} \cot \frac{\pi m}{n} \sin \frac{2\pi km}{n}.$$

H 47. The function $\mu : \mathbf{N} \rightarrow \mathbf{C}$ is defined by

$$\mu(n) = \sum_{k \in R_n} \left(\cos \frac{2k\pi}{n} + i \sin \frac{2k\pi}{n} \right),$$

where $R_n = \{k \in \mathbf{N} | 1 \leq k \leq n, \gcd(k, n) = 1\}$. Show that $\mu(n)$ is an integer for all positive integer n .

10.5. Functional Equations.

H 48. Prove that there is a function f from the set of all natural numbers into itself such that $f(f(n)) = n^2$ for all $n \in \mathbf{N}$.

H 49. Find all surjective function $f : \mathbf{N} \rightarrow \mathbf{N}$ satisfying the condition

$$m|n \iff f(m)|f(n), \quad m, n \in \mathbf{N}.$$

H 50. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(n+1) > f(f(n)), \quad n \in \mathbf{N}.$$

H 51. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(f(f(n))) + f(f(n)) + f(n) = 3n, \quad n \in \mathbf{N}.$$

H 52. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(f(m) + f(n)) = m + n, \quad m, n \in \mathbf{N}.$$

H 53. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f^{(19)}(n) + 97f(n) = 98n + 232, \quad n \in \mathbf{N}.$$

H 54. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(f(n)) + f(n) = 2n + 2001 \text{ or } 2n + 2002, \quad n \in \mathbf{N}.$$

H 55. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(f(f(n))) + 6f(n) = 3f(f(n)) + 4n + 2001, \quad n \in \mathbf{N}.$$

H 56. Find all functions $f : \mathbf{N}_0 \rightarrow \mathbf{N}_0$ such that

$$f(f(n)) + f(n) = 2n + 6, \quad n \in \mathbf{N}_0.$$

H 57. Find all functions $f : \mathbf{N}_0 \rightarrow \mathbf{N}_0$ such that

$$f(m + f(n)) = f(f(m)) + f(n), \quad m, n \in \mathbf{N}_0.$$

H 58. Find all functions $f : \mathbf{N}_0 \rightarrow \mathbf{N}_0$ such that

$$mf(n) + nf(m) = (m+n)f(m^2 + n^2), \quad m, n \in \mathbf{N}_0.$$

H 59. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

- (1) $f(2) = 2$
 (2) $f(mn) = f(m)f(n)$, $m, n \in \mathbf{N}$,
 (3) $f(n+1) > f(n)$, $n \in \mathbf{N}$

H 60. Find all functions $f : \mathbf{Z} \rightarrow \mathbf{Z}$ such that

$$f(f(m)) = m + 1, \quad m \in \mathbf{Z}$$

H 61. Find all functions $f : \mathbf{Z} \rightarrow \mathbf{Z}$ such that

- (1) $f(m+8) \leq f(m) + 8$, $m \in \mathbf{Z}$,
 (2) $f(m+11) \geq f(m) + 11$, $m \in \mathbf{Z}$

H 62. Find all functions $f : \mathbf{Z} \rightarrow \mathbf{Z}$ such that

$$f(m + f(n)) = f(m) - n, \quad m, n \in \mathbf{Z}.$$

H 63. Find all functions $f : \mathbf{Z} \rightarrow \mathbf{Z}$ such that

$$f(m + f(n)) = f(m) + n, \quad m, n \in \mathbf{Z}.$$

H 64. Find all functions $h : \mathbf{Z} \rightarrow \mathbf{Z}$ such that

$$h(x+y) + h(xy) = h(x)h(y) + 1, \quad x, y \in \mathbf{Z}.$$

H 65. Find all functions $f : \mathbb{Q} \rightarrow \mathbf{R}$ such that

$$f(xy) = f(x)f(y) - f(x+y) + 1, \quad x, y \in \mathbb{Q}.$$

H 66. Find all functions $f : \mathbb{Q}^+ \rightarrow \mathbb{Q}^+$ such that

$$f\left(x + \frac{y}{x}\right) = f(x) + \frac{f(y)}{f(x)} + 2y, \quad x, y \in \mathbb{Q}^+.$$

H 67. Find all functions $f : \mathbb{Q} \rightarrow \mathbb{Q}$ such that

$$f(x+y) + f(x-y) = 2(f(x) + f(y)), \quad x, y \in \mathbb{Q}.$$

H 68. Find all functions $f, g, h : \mathbb{Q} \rightarrow \mathbb{Q}$ such that

$$f(x + g(y)) = g(h(f(x))) + y, \quad x, y \in \mathbb{Q}.$$

H 69. Find all functions $f : \mathbb{Q}^+ \rightarrow \mathbb{Q}^+$ such that

- (1) $f(x+1) = f(x) + 1$, $x \in \mathbb{Q}^+$,
 (2) $f(x^2) = f(x)^2$, $x \in \mathbb{Q}^+$.

H 70. Let \mathbb{Q}^+ be the set of positive rational numbers. Construct a function $f : \mathbb{Q}^+ \rightarrow \mathbb{Q}^+$ such that

$$f(xf(y)) = \frac{f(x)}{y}$$

for all $x, y \in \mathbb{Q}^+$.

H 71. A function f is defined on the positive integers by

$$\begin{aligned} f(1) &= 1, & f(3) &= 3, \\ f(2n) &= f(n), \\ f(4n+1) &= 2f(2n+1) - f(n), \\ f(4n+3) &= 3f(2n+1) - 2f(n), \end{aligned}$$

for all positive integers n .

Determine the number of positive integers n , less than or equal to 1988, for which $f(n) = n$.

H 72. Consider all functions f from the set \mathbf{N} of all positive integers into itself satisfying $f(t^2 f(s)) = s(f(t))^2$ for all s and t in \mathbf{N} . Determine the least possible value of $f(1998)$.

H 73. The function $f(n)$ is defined for all positive integers n and takes on non-negative integer values. Also, for all m, n

$$f(m+n) - f(m) - f(n) = 0 \text{ or } 1, \quad f(2) = 0, \quad f(3) > 0, \quad \text{and } f(9999) = 3333.$$

Determine $f(1982)$.

H 74. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(f(m) + f(n)) = m + n, \quad m, n \in \mathbf{N}$$

H 75. Find all surjective functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(n) \geq n + (-1)^n, \quad m, n \in \mathbf{N}$$

H 76. Find all functions $f : \mathbf{Z} - \{0\} \rightarrow \mathbb{Q}$ such that

$$f\left(\frac{x+y}{3}\right) = \frac{f(x) + f(y)}{2}, \quad x, y \in \mathbf{Z} - \{0\}$$

H 77. Find all functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(f(f(n))) + f(f(n)) + f(n) = 3n, \quad n \in \mathbf{N}.$$

H 78. Find all strictly increasing functions $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f(f(n)) = 3n, \quad n \in \mathbf{N}.$$

H 79. Find all functions $f : \mathbf{Z}^2 \rightarrow \mathbf{R}^+$ such that

$$f(i, j) = \frac{f(i+1, j) + f(i, j+1) + f(i-1, j) + f(i, j-1)}{4}, \quad i, j \in \mathbf{Z}.$$

H 80. Find all functions $f : \mathbb{Q} \rightarrow \mathbb{Q}$ such that

$$f(x+y+z) + f(x-y) + f(y-z) + f(z-x) = 3f(x) + 3f(y) + 3f(z), \quad x, y, z \in \mathbb{Q}.$$

H 81. Show that there exists a bijective function $f : \mathbf{N}_0 \rightarrow \mathbf{N}_0$ such that

$$f(3mn + m + n) = 4f(m)f(n) + f(m) + f(n), \quad m, n \in \mathbf{N}_0.$$

H 82. Show that there exists a function $f : \mathbf{N} \rightarrow \mathbf{N}$ such that

$$f^{(1996)}(n) = 2n, \quad n \in \mathbf{N}.$$

11. POLYNOMIALS

The only way to learn Mathematics is to do Mathematics. Paul Halmos

I 1. Suppose $p(x) \in \mathbf{Z}[x]$ and $P(a)P(b) = -(a - b)^2$ for some distinct $a, b \in \mathbf{Z}$. Prove that $P(a) + P(b) = 0$.

I 2. Prove that there is no nonconstant polynomial $f(x)$ with integral coefficients such that $f(n)$ is prime for all $n \in \mathbf{N}$.

I 3. Let $n \geq 2$ be an integer. Prove that if $k^2 + k + n$ is prime for all integers k such that $0 \leq k \leq \sqrt{\frac{n}{3}}$, then $k^2 + k + n$ is prime for all integers k such that $0 \leq k \leq n - 2$.

I 4. A prime p has decimal digits $p_n p_{n-1} \cdots p_0$ with $p_n > 1$. Show that the polynomial $p_n x^n + p_{n-1} x^{n-1} + \cdots + p_1 x + p_0$ cannot be represented as a product of two nonconstant polynomials with integer coefficients

I 5. (Eisenstein's Criterion) Let $f(x) = a_n x^n + \cdots + a_1 x + a_0$ be a nonconstant polynomial with integer coefficients. If there is a prime p such that p divides each of $a_0, a_1, \cdots, a_{n-1}$ but p does not divide a_n and p^2 does not divide a_0 , then $f(x)$ is irreducible in $\mathbb{Q}[x]$.

I 6. Prove that for a prime p , $x^{p-1} + x^{p-2} + \cdots + x + 1$ is irreducible in $\mathbb{Q}[x]$.

I 7. Let $f(x) = x^n + 5x^{n-1} + 3$, where $n > 1$ is an integer. Prove that $f(x)$ cannot be expressed as the product of two nonconstant polynomials with integer coefficients.

I 8. (Eugen Netto) Show that a polynomial of odd degree $2m + 1$ over \mathbf{Z} ,

$$f(x) = c_{2m+1} x^{2m+1} + \cdots + c_1 x + c_0,$$

is irreducible if there exists a prime p such that

$$p \nmid c_{2m+1}, p \mid c_{m+1}, c_{m+2}, \cdots, c_{2m}, p^2 \mid c_0, c_1, \cdots, c_m, \text{ and } p^3 \nmid c_0.$$

I 9. For non-negative integers n and k , let $P_{n,k}(x)$ denote the rational function

$$\frac{(x^n - 1)(x^n - x) \cdots (x^n - x^{k-1})}{(x^k - 1)(x^k - x) \cdots (x^k - x^{k-1})}.$$

Show that $P_{n,k}(x)$ is actually a polynomial for all $n, k \in \mathbf{N}$.

I 10. Suppose that the integers a_1, a_2, \cdots, a_n are distinct. Show that

$$(x - a_1)(x - a_2) \cdots (x - a_n) - 1$$

cannot be expressed as the product of two nonconstant polynomials with integer coefficients.

I 11. Show that the polynomial $x^8 + 98x^4 + 1$ can be expressed as the product of two nonconstant polynomials with integer coefficients.

I 12. Prove that if the integers a_1, a_2, \dots, a_n are all distinct, then the polynomial

$$(x - a_1)^2(x - a_2)^2 \cdots (x - a_n)^2 + 1$$

cannot be expressed as the product of two nonconstant polynomials with integer coefficients.

I 13. On Christmas Eve, 1983, Dean Jixon, the famous seer who had made startling predictions of the events of the preceding year that the volcanic and seismic activities of 1980 and 1981 were connected with mathematics. The diminishing of this geological activity depended upon the existence of an elementary proof of the irreducibility of the polynomial

$$P(x) = x^{1981} + x^{1980} + 12x^2 + 24x + 1983.$$

Is there such a proof?

12. SEQUENCES OF INTEGERS

A peculiarity of the higher arithmetic is the great difficulty which has often been experienced in proving simple general theorems which had been suggested quite naturally by numerical evidence. Harold Davenport

12.1. Linear Recurrences.

J 1. An integer sequence $\{a_n\}_{n \geq 1}$ is defined by

$$a_0 = 0, a_1 = 1, a_{n+2} = 2a_{n+1} + a_n$$

Show that 2^k divides a_n if and only if 2^k divides n .

J 2. The Fibonacci sequence $\{F_n\}$ is defined by

$$F_1 = 1, F_2 = 1, F_{n+2} = F_{n+1} + F_n.$$

Show that $\gcd(F_m, F_n) = F_{\gcd(m,n)}$ for all $m, n \in \mathbf{N}$.

J 3. The Fibonacci sequence $\{F_n\}$ is defined by

$$F_1 = 1, F_2 = 1, F_{n+2} = F_{n+1} + F_n.$$

Show that $F_{mn-1} - F_{n-1}^m$ is divisible by F_n^2 for all $m \geq 1$ and $n > 1$.

J 4. The Fibonacci sequence $\{F_n\}$ is defined by

$$F_1 = 1, F_2 = 1, F_{n+2} = F_{n+1} + F_n.$$

Show that $F_{mn} - F_{n+1}^m + F_{n-1}^m$ is divisible by F_n^3 for all $m \geq 1$ and $n > 1$.

J 5. The Fibonacci sequence $\{F_n\}$ is defined by

$$F_1 = 1, F_2 = 1, F_{n+2} = F_{n+1} + F_n.$$

Show that $F_{2n-1}^2 + F_{2n+1}^2 + 1 = 3F_{2n-1}F_{2n+1}$ for all $n \geq 1$.

J 6. Prove that no Fibonacci number can be factored into a product of two smaller Fibonacci numbers, each greater than 1.

J 7. Let m be a positive integer. Define the sequence $\{a_n\}_{n \geq 0}$ by

$$a_0 = 0, a_1 = m, a_{n+1} = m^2 a_n - a_{n-1}.$$

Prove that an ordered pair (a, b) of non-negative integers, with $a \leq b$, gives a solution to the equation

$$\frac{a^2 + b^2}{ab + 1} = m^2$$

if and only if (a, b) is of the form (a_n, a_{n+1}) for some $n \geq 0$.

J 8. Let x_n and y_n be two sequences defined recursively as follows

$$x_0 = 1, x_1 = 4, x_{n+2} = 3x_{n+1} - x_n$$

$$y_0 = 1, y_1 = 2, y_{n+2} = 3y_{n+1} - y_n$$

for all $n = 0, 1, 2, \dots$.

a) Prove that $x_n^2 - 5y_n^2 + 4 = 0$ for all non-negative integers.

b) Suppose that a, b are two positive integers such that $a^2 - 5b^2 + 4 = 0$. Prove that there exists a non-negative integer k such that $a = x_k$ and $b = y_k$.

J 9. Let $\{u_n\}_{n \geq 0}$ be a sequence of positive integers defined by

$$u_0 = 1, u_{n+1} = au_n + b,$$

where $a, b \in \mathbf{N}$. Prove that for any choice of a and b , the sequence $\{u_n\}_{n \geq 0}$ contains infinitely many composite numbers.

J 10. The sequence $\{y_n\}_{n \geq 1}$ is defined by

$$y_1 = y_2 = 1, y_{n+2} = (4k - 5)y_{n+1} - y_n + 4 - 2k \quad (n \in \mathbf{N}).$$

Determine all integers k such that each term of this sequence is a perfect square.

J 11. Let the sequence $\{K_n\}_{n \geq 1}$ be defined by

$$K_1 = 2, K_2 = 8, K_{n+2} = 3K_{n+1} - K_n + 5(-1)^n.$$

Prove that if K_n is prime, then n must be a power of 3.

J 12. The sequence $\{a_n\}_{n \geq 1}$ is defined by

$$a_1 = 1, a_2 = 12, a_3 = 20, a_{n+3} = 2a_{n+2} + 2a_{n+1} - a_n \quad (n \in \mathbf{N}).$$

Prove that $1 + 4a_n a_{n+1}$ is a square for all $n \in \mathbf{N}$.

J 13. The sequence $\{x_n\}_{n \geq 1}$ is defined by

$$x_1 = x_2 = 1, x_{n+2} = 14x_{n+1} - x_n - 4 \quad (n = 1, 2, \dots)$$

Prove that x_n is always a perfect square.

12.2. Recursive Sequences.

J 14. Let $P(x)$ be a nonzero polynomial with integral coefficients. Let $a_0 = 0$ and for $i \geq 0$ define $a_{i+1} = P(a_i)$. Show that $\gcd(a_m, a_n) = a_{\gcd(m,n)}$ for all $m, n \in \mathbf{N}$

J 15. An integer sequence $\{a_n\}_{n \geq 1}$ is defined by

$$a_1 = 1, a_{n+1} = a_n + \lfloor \sqrt{a_n} \rfloor$$

Show that a_n is a square if and only if $n = 2^k + k - 2$ for some $k \in \mathbf{N}$.

J 16. Let $f(n) = n + \lfloor \sqrt{n} \rfloor$. Prove that, for every positive integer m , the sequence

$$m, f(m), f(f(m)), f(f(f(m))), \dots$$

contains at least one square of an integer.

J 17. The sequence $\{a_n\}_{n \geq 1}$ is defined by

$$a_1 = 1, a_2 = 2, a_3 = 24, a_{n+2} = \frac{6a_{n-1}^2 a_{n-3} - 8a_{n-1} a_{n-2}^2}{a_{n-2} a_{n-3}} \quad (n \geq 4)$$

Show that a_n is an integer for all n .

J 18. Show that there is a unique sequence of integers $\{a_n\}_{n \geq 1}$ with

$$a_1 = 1, a_2 = 2, a_4 = 12, a_{n+1} a_{n-1} = a_n^2 + 1 \quad (n \geq 2).$$

J 19. The sequence $\{a_n\}_{n \geq 1}$ is defined by

$$a_1 = 1, a_{n+1} = 2a_n + \sqrt{3a_n^2 + 1} \quad (n \geq 1)$$

Show that a_n is an integer for every n .

J 20. Prove that the sequence $\{y_n\}_{n \geq 1}$ defined by

$$y_0 = 1, y_{n+1} = \frac{1}{2} \left(3y_n + \sqrt{5a_n^2 - 4} \right) \quad (n \geq 0)$$

consists only of integers.

J 21. (C. von Staudt) The Bernoulli sequence¹⁶ $\{B_n\}_{n \geq 0}$ is defined by

$$B_0 = 1, B_n = -\frac{1}{n+1} \sum_{k=0}^n \binom{n+1}{k} B_k \quad (n \geq 1)$$

Show that for all $n \in \mathbf{N}$,

$$(-1)^n B_n - \sum \frac{1}{p},$$

is an integer where the summation being extended over the primes p such that $p|2k-1$.

J 22. An integer sequence $\{a_n\}_{n \geq 1}$ is defined by

$$a_1 = 2, a_{n+1} = \left\lfloor \frac{3}{2} a_n \right\rfloor$$

Show that it has infinitely many even and infinitely many odd integers.

J 23. An integer sequence satisfies $a_{n+1} = a_n^3 + 1999$. Show that it contains at most one square.

J 24. Let $a_1 = 11^{11}$, $a_2 = 12^{12}$, $a_3 = 13^{13}$, and

$$a_n = |a_{n-1} - a_{n-2}| + |a_{n-2} - a_{n-3}|, n \geq 4.$$

Determine $a_{14^{14}}$.

J 25. Let k be a fixed positive integer. The infinite sequence a_n is defined by the formulae

$$a_1 = k + 1, a_{n+1} = a_n^2 - k a_n + k \quad (n \geq 1).$$

Show that if $m \neq n$, then the numbers a_m and a_n are relatively prime.

¹⁶ $B_0 = 1, B_1 = -\frac{1}{2}, B_2 = \frac{1}{6}, B_3 = 0, B_4 = -\frac{1}{30}, B_5 = 0, B_6 = \frac{1}{42}, \dots$

J 26. The sequence $\{x_n\}$ is defined by

$$x_0 \in [0, 1], \quad x_{n+1} = 1 - |1 - 2x_n|.$$

Prove that the sequence is periodic if and only if x_0 is irrational.

J 27. Let x_1 and x_2 be relatively prime positive integers. For $n \geq 2$, define $x_{n+1} = x_n x_{n-1} + 1$.

(a) Prove that for every $i > 1$, there exists $j > i$ such that x_i^i divides x_j^j .

(b) Is it true that x_1 must divide x_j^j for some $j > 1$?

J 28. For a given positive integer k denote the square of the sum of its digits by $f_1(k)$ and let $f_{n+1}(k) = f_1(f_n(k))$. Determine the value of $f_{1991}(2^{1990})$.

J 29. Define a sequence $\{a_i\}$ by $a_1 = 3$ and $a_{i+1} = 3^{a_i}$ for $i \geq 1$. Which integers between 00 and 99 inclusive occur as the last two digits in the decimal expansion of infinitely many a_i ?

J 30. A sequence of integers, $\{a_n\}_{n \geq 1}$ with $a_1 > 0$, is defined by

$$\begin{aligned} a_{n+1} &= \frac{a_n}{2} && \text{if } n \equiv 0 \pmod{4}, \\ a_{n+1} &= 3a_n + 1 && \text{if } n \equiv 1 \pmod{4}, \\ a_{n+1} &= 2a_n - 1 && \text{if } n \equiv 2 \pmod{4}, \\ a_{n+1} &= \frac{a_n + 1}{4} && \text{if } n \equiv 3 \pmod{4}. \end{aligned}$$

Prove that there is an integer m such that $a_m = 1$.

J 31. Given is an integer sequence $\{a_n\}_{n \geq 0}$ such that $a_0 = 2$, $a_1 = 3$ and, for all positive integers $n \geq 1$, $a_{n+1} = 2a_{n-1}$ or $a_{n+1} = 3a_n - 2a_{n-1}$. Does there exist a positive integer k such that $1600 < a_k < 2000$?

J 32. A sequence with first two terms equal 1 and 24 respectively is defined by the following rule: each subsequent term is equal to the smallest positive integer which has not yet occurred in the sequence and is not coprime with the previous term. Prove that all positive integers occur in this sequence.

J 33. Each term of a sequence of natural numbers is obtained from the previous term by adding to it its largest digit. What is the maximal number of successive odd terms in such a sequence?

J 34. In the sequence $1, 0, 1, 0, 1, 0, 3, 5, \dots$, each member after the sixth one is equal to the last digit of the sum of the six members just preceding it. Prove that in this sequence one cannot find the following group of six consecutive members :

$$0, 1, 0, 1, 0, 1$$

J 35. Let a , and b be odd positive integers. Define the sequence (f_n) by putting $f_1 = a$, $f_2 = b$, and by letting f_n for $n \geq 3$ be the greatest odd divisor of $f_{n-1} + f_{n-2}$. Show that f_n is constant for sufficiently large n and determine the eventual value as a function of a and b .

J 36. Numbers $d(n, m)$ with m, n integers, $0 \leq m \leq n$, are defined by $d(n, 0) = d(n, n) = 1$ ($n \geq 0$), $md(n, m) = md(n-1, m) + (2n-m)d(n-1, m-1)$ ($0 < m < n$). Prove that $d(n, m)$ are integers for all $m, n \in \mathbf{N}$.

J 37. Let k be a given positive integer. The sequence x_n is defined as follows: $x_1 = 1$ and x_{n+1} is the least positive integer which is not in $\{x_1, x_2, \dots, x_n, x_1+k, x_2+2k, \dots, x_n+nk\}$. Show that there exist real number a such that $x_n = [an]$ for all positive integer n .

J 38. Let $\{a_n\}_{n \geq 1}$ be a sequence of positive integers such that

$$0 < a_{n+1} - a_n \leq 2001 \text{ for all } n \in \mathbf{N}.$$

Show that there are infinitely many pairs (p, q) of positive integers such that $p > q$ and $a_q \mid a_p$.

J 39. Let p be an odd prime p such that $2h \not\equiv 1 \pmod{p}$ for all $h \in \mathbf{N}$ with $h < p-1$, and let a be an even integer with $a \in (\frac{p}{2}, p)$. The sequence $\{a_n\}_{n \geq 0}$ is defined by $a_0 = a$, $a_{n+1} = p - b_n$ ($n \geq 0$), where b_n is the greatest odd divisor of a_n . Show that the sequence $\{a_n\}_{n \geq 0}$ is periodic and find its minimal (positive) period.

J 40. Let $p \geq 3$ be a prime number. The sequence $\{a_n\}_{n \geq 1}$ is defined by $a_n = n$ for all $0 \leq n \leq p-1$, and $a_n = a_{n-1} + a_{n-p}$, for all $n \geq p$. Compute $a_{p^3} \pmod{p}$.

J 41. Let $\{u_n\}_{n \geq 0}$ be a sequence of integers satisfying the recurrence relation $u_{n+2} = u_{n+1}^2 - u_n$ ($n \in \mathbf{N}$). Suppose that $u_0 = 39$ and $u_1 = 45$. Prove that 1986 divides infinitely many terms of this sequence.

J 42. The sequence $\{a_n\}_{n \geq 1}$ is defined by $a_1 = 1$ and

$$a_{n+1} = \frac{a_n}{2} + \frac{1}{4a_n} \quad (n \in \mathbf{N}).$$

Prove that $\sqrt{\frac{2}{2a_n^2-1}}$ is a positive integer for $n > 1$.

J 43. Let k be a positive integer. Prove that there exists an infinite monotone increasing sequence of integers $\{a_n\}_{n \geq 1}$ such that

$$a_n \text{ divides } a_{n+1}^2 + k \quad \text{and} \quad a_{n+1} \text{ divides } a_n^2 + k$$

for all $n \in \mathbf{N}$.

J 44. Each term of an infinite sequence of natural numbers is obtained from the previous term by adding to it one of its nonzero digits. Prove that this sequence contains an even number.

J 45. In an increasing infinite sequence of positive integers, every term starting from the 2002-th term divides the sum of all preceding terms. Prove that every term starting from some term is equal to the sum of all preceding terms.

J 46. The sequence $\{x_n\}_{n \geq 1}$ is defined by

$$x_1 = 2, x_{n+1} = \frac{2 + x_n}{1 - 2x_n} \quad (n \in \mathbf{N}).$$

Prove that (a) $x_n \neq 0$ for all $n \in \mathbf{N}$ and (b) $\{x_n\}_{n \geq 1}$ is not periodic.

J 47. (A. Perlin) The sequence of integers $\{x_n\}$ is defined as follows :

$$x_1 = 1, \quad x_{n+1} = 1 + x_1^2 + \cdots + x_n^2 \quad (n = 1, 2, 3, \dots).$$

Prove that there are no squares of natural numbers in this sequence except x_1 .

J 48. The first four terms of an infinite sequence S of decimal digits are 1, 9, 8, 2, and succeeding terms are given by the final digit in the sum of the four immediately preceding terms. Thus S begins 1, 9, 8, 2, 0, 9, 9, 0, 8, 6, 3, 7, 4, \dots . Do the digits 3, 0, 4, 4 ever come up consecutively in S ?

12.3. More Sequences.

J 49. Show that the sequence $\{a_n\}_{n \geq 1}$ defined by $a_n = [n\sqrt{2}]$ contains an infinite number of integer powers of 2.

J 50. Let a_n be the last nonzero digit in the decimal representation of the number $n!$. Does the sequence a_1, a_2, a_3, \dots become periodic after a finite number of terms ?

J 51. Let $n > 6$ be an integer and a_1, a_2, \dots, a_k be all the natural numbers less than n and relatively prime to n . If

$$a_2 - a_1 = a_3 - a_2 = \cdots = a_k - a_{k-1} > 0,$$

prove that n must be either a prime number or a power of 2.

J 52. Show that if an infinite arithmetic progression of positive integers contains a square and a cube, it must contain a sixth power.

J 53. Prove that there exist two strictly increasing sequences a_n and b_n such that $a_n(a_n + 1)$ divides $b_n^2 + 1$ for every natural n .

J 54. Let $\{a_n\}$ be a strictly increasing positive integers sequence such that $\gcd(a_i, a_j) = 1$ and $a_{i+2} - a_{i+1} > a_{i+1} - a_i$. Show that the infinite series

$$\sum_{i=1}^{\infty} \frac{1}{a_i}$$

converges.

J 55. Let $\{n_k\}_{k \geq 1}$ be a sequence of natural numbers such that for $i < j$, the decimal representation of n_i does not occur as the leftmost digits of the decimal representation of n_j . Prove that

$$\sum_{k=1}^{\infty} \frac{1}{n_k} \leq \frac{1}{1} + \frac{1}{2} + \cdots + \frac{1}{9}.$$

J 56. An integer sequence $\{a_n\}_{n \geq 1}$ is given such that

$$2^n = \sum_{d|n} a_d$$

for all $n \in \mathbf{N}$. Show that a_n is divisible by n for all $n \in \mathbf{N}$.

J 57. Let q_0, q_1, \dots be a sequence of integers such that

- (i) for any $m > n$, $m - n$ is a factor of $q_m - q_n$, and
- (ii) $|q_n| \leq n^{10}$ for all integers $n \geq 0$.

Show that there exists a polynomial $Q(x)$ satisfying $q_n = Q(n)$ for all n .

J 58. Let a, b be integers greater than 2. Prove that there exists a positive integer k and a finite sequence n_1, n_2, \dots, n_k of positive integers such that $n_1 = a$, $n_k = b$, and $n_i n_{i+1}$ is divisible by $n_i + n_{i+1}$ for each i ($1 \leq i < k$).

J 59. The infinite sequence of 2's and 3's

$$\begin{aligned} &2, 3, 3, 2, 3, 3, 3, 2, 3, 3, 3, 2, 3, 3, 2, 3, 3, \\ &3, 2, 3, 3, 3, 2, 3, 3, 3, 2, 3, 3, 2, 3, 3, 3, 2, \dots \end{aligned}$$

has the property that, if one forms a second sequence that records the number of 3's between successive 2's, the result is identical to the given sequence. Show that there exists a real number r such that, for any n , the n th term of the sequence is 2 if and only if $n = 1 + \lfloor rm \rfloor$ for some nonnegative integer m . (Note: $\lfloor x \rfloor$ denotes the largest integer less than or equal to x .)

J 60. The sequence $\{a_n\}_{n \geq 1}$ is defined by

$$a_n = 1 + 2^2 + 3^3 + \cdots + n^n.$$

Prove that there are infinitely many n such that a_n is composite.

J 61. One member of an infinite arithmetic sequence in the set of natural numbers is a perfect square. Show that there are infinitely many members of this sequence having this property.

J 62. In the sequence 00, 01, 02, 03, \dots , 99 the terms are rearranged so that each term is obtained from the previous one by increasing or decreasing one of its digits by 1 (for example, 29 can be followed by 19, 39, or 28, but not by 30 or 20). What is the maximal number of terms that could remain on their places?

J 63. Does there exist positive integers $a_1 < a_2 < \cdots < a_{100}$ such that for $2 \leq k \leq 100$, the least common multiple of a_{k-1} and a_k is greater than the least common multiple of a_k and a_{k+1} ?

J 64. Does there exist positive integers $a_1 < a_2 < \cdots < a_{100}$ such that for $2 \leq k \leq 100$, the greatest common divisor of a_{k-1} and a_k is greater than the greatest common divisor of a_k and a_{k+1} ?

J 65. Suppose that a and b are distinct real numbers such that

$$a - b, a^2 - b^2, \dots, a^k - b^k, \dots$$

are all integers. Show that a and b are integers.

13. COMBINATORIAL NUMBER THEORY

In great mathematics there is a very high degree of unexpectedness, combined with inevitability and economy. Godfrey Harold Hardy

K 1. (Erdős) Suppose all the pairs of a positive integers from a finite collection

$$A = \{a_1, a_2, \dots\}$$

are added together to form a new collection

$$A^* = \{a_i + a_j \mid 1 \leq i < j \leq n\}.$$

For example, $A = \{2, 3, 4, 7\}$ would yield $A^* = \{5, 6, 7, 9, 10, 11\}$ and $B = \{1, 4, 5, 6\}$ would give $B^* = \{5, 6, 7, 9, 10, 11\}$. These examples show that it's possible for different collections A and B to generate the same collections A^* and B^* . Show that if $A^* = B^*$ for different sets A and B , then $|A| = |B|$ and $|A| = |B|$ must be a power of 2.

K 2. Let p be a prime. Find all positive integers k such that the set $\{1, 2, \dots, k\}$ can be partitioned into p subsets with equal sum of elements.

K 3. Prove that the set of integers of the form $2^k - 3$ ($k = 2, 3, \dots$) contains an infinite subset in which every two members are relatively prime.

K 4. The set of positive integers is partitioned into finitely many subsets. Show that some subset S has the following property : for every positive integer n , S contains infinitely many multiples of n .

K 5. Let M be a positive integer and consider the set

$$S = \{n \in \mathbf{N} \mid M^2 \leq n < (M + 1)^2\}.$$

Prove that the products of the form ab with $a, b \in S$ are distinct.

K 6. Let S be a set of integers such that

- there exist $a, b \in S$ with $\gcd(a, b) = \gcd(a - 2, b - 2) = 1$.
- if x and y are elements of S , then $x^2 - y$ also belongs to S .

Prove that S is the set of all integers.

K 7. Show that for each $n \geq 2$, there is a set S of n integers such that $(a - b)^2$ divides ab for every distinct $a, b \in S$

K 8. Let a and b be positive integers greater than 2. Prove that there exists a positive integer k and a finite sequence n_1, \dots, n_k of positive integers such that $n_1 = a$, $n_k = b$, and $n_i n_{i+1}$ is divisible by $n_i + n_{i+1}$ for each i ($1 \leq i \leq k$).

K 9. Let n be an integer, and let X be a set of $n+2$ integers each of absolute value at most n . Show that there exist three distinct numbers $a, b, c \in X$ such that $c = a + b$.

K 10. Let $m \geq 2$ be an integer. Find the smallest integer $n > m$ such that for any partition of the set $\{m, m+1, \dots, n\}$ into two subsets, at least one subset contains three numbers a, b, c such that $c = a^b$.

K 11. Let $S = \{1, 2, 3, \dots, 280\}$. Find the smallest integer n such that each n -element subset of S contains five numbers which are pairwise relatively prime.

K 12. Let m and n be positive integers. If x_1, x_2, \dots, x_m are positive integers whose average is less than $n+1$ and if y_1, y_2, \dots, y_n are positive integers whose average is less than $m+1$, prove that some sum of one or more x 's equals some sum of one or more y 's.

K 13. Let n and k be given relatively prime natural numbers, $k < n$. Each number in the set $M = \{1, 2, \dots, n-1\}$ is colored either blue or white. It is given that

- for each $i \in M$, both i and $n-i$ have the same color;
- for each $i \in M, i \neq k$, both i and $|i-k|$ have the same color.

Prove that all numbers in M have the same color.

K 14. Let p be a prime number, $p \geq 5$, and k be a digit in the p -adic representation of positive integers. Find the maximal length of a non constant arithmetic progression whose terms do not contain the digit k in their p -adic representation.

K 15. Is it possible to choose 1983 distinct positive integers, all less than or equal to 10^5 , no three of which are consecutive terms of an arithmetic progression?

K 16. Is it possible to find 100 positive integers not exceeding 25000 such that all pairwise sums of them are different?

K 17. Find the maximum number of pairwise disjoint sets of the form

$$S_{a,b} = \{n^2 + an + b \mid n \in \mathbf{Z}\},$$

with $a, b \in \mathbf{Z}$.

K 18. Let p be an odd prime number. How many p -element subsets A of $\{1, 2, \dots, 2p\}$ are there, the sum of whose elements is divisible by p ?

K 19. Let $m, n \geq 2$ be positive integers, and let a_1, a_2, \dots, a_n be integers, none of which is a multiple of m^{n-1} . Show that there exist integers e_1, e_2, \dots, e_n , not all zero, with $|e_i| < m$ for all i , such that $e_1 a_1 + e_2 a_2 + \dots + e_n a_n$ is a multiple of m^n .

K 20. Determine the smallest integer $n \geq 4$ for which one can choose four different numbers a, b, c , and d from any n distinct integers such that $a + b - c - d$ is divisible by 20

K 21. A sequence of integers a_1, a_2, a_3, \dots is defined as follows : $a_1 = 1$, and for $n \geq 1$, a_{n+1} is the smallest integer greater than a_n such that $a_i + a_j \neq 3a_k$ for any i, j , and k in $\{1, 2, 3, \dots, n+1\}$, not necessarily distinct. Determine a_{1998} .

K 22. Prove that for each positive integer n , there exists a positive integer with the following properties :

- It has exactly n digits.
- None of the digits is 0.
- It is divisible by the sum of its digits.

K 23. Let k, m, n be integers such that $1 < n \leq m - 1 \leq k$. Determine the maximum size of a subset S of the set $\{1, 2, \dots, k\}$ such that no n distinct elements of S add up to m .

K 24. Find the number of subsets of $\{1, 2, \dots, 2000\}$, the sum of whose elements is divisible by 5.

K 25. Let A be a non-empty set of positive integers. Suppose that there are positive integers b_1, \dots, b_n and c_1, \dots, c_n such that

- (i) for each i the set $b_i A + c_i = \{b_i a + c_i \mid a \in A\}$ is a subset of A , and
- (ii) the sets $b_i A + c_i$ and $b_j A + c_j$ are disjoint whenever $i \neq j$.

Prove that

$$\frac{1}{b_1} + \dots + \frac{1}{b_n} \leq 1.$$

K 26. A set of three nonnegative integers $\{x, y, z\}$ with $x < y < z$ is called historic if $\{z - y, y - x\} = \{1776, 2001\}$. Show that the set of all nonnegative integers can be written as the unions of pairwise disjoint historic sets.

K 27. Let p and q be relatively prime positive integers. A subset S of $\{0, 1, 2, \dots\}$ is called ideal if $0 \in S$ and, for each element $n \in S$, the integers $n + p$ and $n + q$ belong to S . Determine the number of ideal subsets of $\{0, 1, 2, \dots\}$.

K 28. Prove that the set of positive integers cannot be partitioned into three nonempty subsets such that, for any two integers x, y taken from two different subsets, the number $x^2 - xy + y^2$ belongs to the third subset.

K 29. Let A be a set of N residues (mod N^2). Prove that there exists a set B of N residues (mod N^2) such that the set $A + B = \{a + b \mid a \in A, b \in B\}$ contains at least half of all the residues (mod N^2).

K 30. Determine the largest positive integer n for which there exists a set S with exactly n numbers such that

- (i) each member in S is a positive integer not exceeding 2002,
- (ii) if a and b are two (not necessarily different) numbers in S , then their product ab does not belong to S .

K 31. Prove that, for any integer $a_1 > 1$, there exist an increasing sequence of positive integers a_1, a_2, a_3, \dots such that

$$a_1 + a_2 + \dots + a_n \mid a_1^2 + a_2^2 + \dots + a_n^2$$

for all $k \in \mathbf{N}$.

K 32. An odd integer $n \geq 3$ is said to be "nice" if and only if there is at least one permutation a_1, \dots, a_n of $1, \dots, n$ such that the n sums $a_1 - a_2 + a_3 - \dots - a_{n-1} + a_n, a_2 - a_3 + a_3 - \dots - a_n + a_1, a_3 - a_4 + a_5 - \dots - a_1 + a_2, \dots, a_n - a_1 + a_2 - \dots - a_{n-2} + a_{n-1}$ are all positive. Determine the set of all "nice" integers.

K 33. Assume that the set of all positive integers is decomposed into r distinct subsets A_1, A_2, \dots, A_r $A_1 \cup A_2 \cup \dots \cup A_r = \mathbf{N}$. Prove that one of them, say A_i , has the following property : There exist a positive integer m such that for any k one can find numbers a_1, \dots, a_k in A_i with $0 < a_{j+1} - a_j \leq m$ ($1 \leq j \leq k - 1$).

K 34. Determine for which positive integers k , the set

$$X = \{1990, 1990 + 1, 1990 + 2, \dots, 1990 + k\}$$

can be partitioned into two disjoint subsets A and B such that the sum of the elements of A is equal to the sum of the elements of B .

K 35. Prove that $n \geq 3$ be a prime number and $a_1 < a_2 < \dots < a_n$ be integers. Prove that a_1, \dots, a_n is an arithmetic progression if and only if there exists a partition of $\{0, 1, 2, \dots\}$ into classes A_1, A_2, \dots, A_n such that

$$a_1 + A_1 = a_2 + A_2 = \dots = a_n + A_n,$$

where $x + A$ denotes the set $\{x + a \mid a \in A\}$.

K 36. Let a and b be non-negative integers such that $ab \geq c^2$ where c is an integer. Prove that there is a positive integer n and integers $x_1, x_2, \dots, x_n, y_1, y_2, \dots, y_n$ such that

$$x_1^2 + \dots + x_n^2 = a, y_1^2 + \dots + y_n^2 = b, x_1y_1 + \dots + x_ny_n = c$$

K 37. Let n, k be positive integers such that n is not divisible by 3 and k is greater or equal to n . Prove that there exists a positive integer m which is divisible by n and the sum of its digits in the decimal representation is k .

K 38. Prove that for every real number M there exists an infinite arithmetical progression such that

- each term is a positive integer and the common difference is not divisible by 10.
- the sum of digits of each term exceeds M .

K 39. Find the smallest positive integer n , for which there exist n different positive integers a_1, a_2, \dots, a_n satisfying the conditions :

- a) the smallest common multiple of a_1, a_2, \dots, a_n is 1985;
 b) for each $i, j \in \{1, 2, \dots, n\}$, the numbers a_i and a_j have a common divisor;
 c) the product $a_1 a_2 \cdots a_n$ is a perfect square and is divisible by 243.

Find all n -tuples (a_1, \dots, a_n) , satisfying a), b), and c).

K 40. Let X be a non-empty set of positive integers which satisfies the following :

- (a) If $x \in X$, then $4x \in X$.
 (b) If $x \in X$, then $\lfloor \sqrt{x} \rfloor \in X$.

Prove that $X = \mathbf{N}$.

K 41. Prove that for every positive integer n there exists an n -digit number divisible by 5^n all of whose digits are odd.

K 42. Let N_n denote the number of ordered n -tuples of positive integers (a_1, a_2, \dots, a_n) such that

$$1/a_1 + 1/a_2 + \dots + 1/a_n = 1.$$

Determine whether N_{10} is even or odd.

K 43. Is it possible to find a set A of eleven positive integers such that no six elements of A have a sum which is divisible by 6 ?

K 44. A set C of positive integers is called good if for every integer k there exist distinct $a, b \in C$ such that the numbers $a+k$ and $b+k$ are not relatively prime. Prove that if the sum of the elements of a good set C equals 2003, then there exists $c \in C$ such that the set $C - \{c\}$ is good.

K 45. Find the set of all positive integers n with the property that the set

$$\{n, n+1, n+2, n+3, n+4, n+5\}$$

can be partitioned into two sets such that the product of the numbers in one set equals the product of the numbers in the other set.

K 46. Suppose p is a prime with $p \equiv 3 \pmod{4}$. Show that for any set of $p-1$ consecutive integers, the set cannot be divided two subsets so that the product of the members of the one set is equal to the product of the members of the other set.

K 47. Let S be the set of all composite positive odd integers less than 79.

(a) Show that S may be written as the union of three (not necessarily disjoint) arithmetic progressions.

(b) Show that S cannot be written as the union of two arithmetic progressions.

K 48. Let a_1, \dots, a_{44} be natural numbers such that

$$0 < a_1 < a_2 < \dots < a_{44} < 125.$$

Prove that at least one of the 43 differences $d_j = a_{j+1} - a_j$ occurs at least 10 times.

K 49. (D. Fomin) Consider the set of all five-digit numbers whose decimal representation is a permutation of the digits 1, 2, 3, 4, 5. Prove that this set can be divided into two groups, in such a way that the sum of the squares of the numbers in each group is the same.

K 50. What's the largest number of elements that a set of positive integers between 1 and 100 inclusive can have if it has the property that none of them is divisible by another ?

K 51. Prove that among 16 consecutive integers it is always possible to find one which is relatively prime to all the rest.

K 52. Is there a set S of positive integers such that a number is in S if and only if it is the sum of two distinct members of S or a sum of two distinct positive integers not in S ?

K 53. Suppose that the set $M = \{1, 2, \dots, n\}$ is split into t disjoint subsets M_1, \dots, M_t where the cardinality of M_i is m_i , and $m_i \geq m_{i+1}$, for $i = 1, \dots, t-1$. Show that if $n > t!e$ then at least one class M_z contains three elements x_i, x_j, x_k with the property that $x_i - x_j = x_k$.

K 54. Let S be a subset of $\{1, 2, 3, \dots, 1989\}$ in which no two members differ by exactly 4 or by exactly 7. What is the largest number of elements S can have ?

K 55. The set M consists of integers, the smallest of which is 1 and the greatest 100. Each member of M , except 1, is the sum of two (possibly identical) numbers in M . Of all such sets, find one with the smallest possible number of elements.

K 56. Show that it is possible to color the set of integers

$$M = \{1, 2, 3, \dots, 1987\},$$

using four colors, so that no arithmetic progression with 10 terms has all its members the same color.

K 57. Prove that every selection of 1325 integers from $M = \{1, 2, \dots, 1987\}$ must contain some three numbers $\{a, b, c\}$ which are relatively prime in pairs, but that can be avoided if only 1324 integers are selected.

K 58. Prove that every infinite sequence S of distinct positive integers contains either

(a) an infinite subsequence such that, for every pair of terms, neither term ever divides the other, or

(b) an infinite subsequence such that, in every pair of terms, one always divides the other.

K 59. *Let $a_1 < a_2 < a_3 < \dots$ be an infinite increasing sequence of positive integers in the number of prime factors of each term, counting repeated factors, is never more than 1987. Prove that it is always possible to extract from A an infinite subsequence $b_1 < b_2 < b_3 < \dots$ such that the greatest common divisor (b_i, b_j) is the same number for every pair of its terms.*

14. ADDITIVE NUMBER THEORY

On Ramanujan, G. H. Hardy Said : I remember once going to see him when he was lying ill at Putney. I had ridden in taxi cab number 1729 and remarked that the number seemed to me rather a dull one, and that I hoped it was not an unfavorable omen. "No," he replied,

"it is a very interesting number; it is the smallest number expressible as the sum of two cubes in two different ways."

L 1. Show that any integer can be expressed as a sum of two squares and a cube.

L 2. Show that each integer n can be written as the sum of five perfect cubes (not necessarily positive).

L 3. Prove that infinitely many positive integers cannot be written in the form

$$x_1^3 + x_2^5 + x_3^7 + x_4^9 + x_5^{11},$$

where $x_1, x_2, x_3, x_4, x_5 \in \mathbf{N}$.

L 4. Determine all positive integers that are expressible in the form

$$a^2 + b^2 + c^2 + c,$$

where a, b, c are integers.

L 5. Show that any positive rational number can be represented as the sum of three positive rational cubes.

L 6. A positive integer n is a square-free integer if there is no prime p such that $p^2 \mid n$. Show that every integer greater than 1 can be written as a sum of two square-free integers.

L 7. Prove that every integer $n \geq 12$ is the sum of two composite numbers.

L 8. Prove that any positive integer can be represented as an aggregate of different powers of 3, the terms in the aggregate being combined by the signs $+$ and $-$ appropriately chosen.

L 9. The integer 9 can be written as a sum of two consecutive integers : $9=4+5$; moreover it can be written as a sum of (more than one) consecutive positive integers in exactly two ways, namely $9=4+5= 2+3+4$. Is there an integer which can be written as a sum of 1990 consecutive integers and which can be written as a sum of (more than one) consecutive integers in exactly 1990 ways ?

L 10. For each positive integer n , $S(n)$ is defined to be the greatest integer such that, for every positive integer $k \leq S(n)$, n^2 can be written as the sum of k positive squares.

- (a) Prove that $S(n) \leq n^2 - 14$ for each $n \geq 4$.
 (b) Find an integer n such that $S(n) = n^2 - 14$.
 (c) Prove that there are infinitely many integers n such that $S(n) = n^2 - 14$.

L 11. For each positive integer n , let $f(n)$ denote the number of ways of representing n as a sum of powers of 2 with nonnegative integer exponents. Representations which differ only in the ordering of their summands are considered to be the same. For instance, $f(4) = 4$, because the number 4 can be represented in the following four ways:

$$4; 2 + 2; 2 + 1 + 1; 1 + 1 + 1 + 1.$$

Prove that, for any integer $n \geq 3$,

$$2^{n^2/4} < f(2^n) < 2^{n^2/2}.$$

L 12. The positive function $p(n)$ is defined as the number of ways that the positive integer n can be written as a sum of positive integers.¹⁷ Show that, for all positive integers $n \geq 2$,

$$2^{\lfloor \sqrt{n} \rfloor} < p(n) < n^{3\lfloor \sqrt{n} \rfloor}.$$

L 13. Let $a_1 = 1, a_2 = 2, a_3, a_4, \dots$ be the sequence of positive integers of the form $2^\alpha 3^\beta$, where α and β are nonnegative integers. Prove that every positive integer is expressible in the form

$$a_{i_1} + a_{i_2} + \dots + a_{i_n},$$

where no summand is a multiple of any other.

L 14. Let n be a non-negative integer. Find non-negative integers a, b, c, d such that

$$a^2 + b^2 + c^2 + d^2 = 7 \cdot 4^n.$$

L 15. Find all integers $m > 1$ such that m^3 is a sum of m squares of consecutive integers.

L 16. Prove that there exist infinitely many integers n such that $n, n+1, n+2$ are each the sum of the squares of two integers.

L 17. (Jacobsthal) Let p be a prime number of the form $4k + 1$. Suppose that r is a quadratic residue of p and that s is a quadratic nonresidue of p . Show that $p = a^2 + b^2$, where

$$a = \frac{1}{2} \sum_{i=1}^{p-1} \left(\frac{i(i^2 - r)}{p} \right), b = \frac{1}{2} \sum_{i=1}^{p-1} \left(\frac{i(i^2 - s)}{p} \right).$$

Here, $\left(\frac{k}{p} \right)$ denotes the Legendre Symbol.

¹⁷For example, $5 = 4 + 1 = 3 + 2 = 3 + 1 + 1 = 2 + 2 + 1 = 2 + 1 + 1 + 1 = 1 + 1 + 1 + 1 + 1$ gives us $p(5) = 7$.

L 18. Let p be a prime with $p \equiv 1 \pmod{4}$. Let a be the unique integer such that

$$p = a^2 + b^2, \quad a \equiv -1 \pmod{4}, \quad b \equiv 0 \pmod{2}$$

Prove that

$$\sum_{i=0}^{p-1} \left(\frac{i^3 + 6i^2 + i}{p} \right) = 2 \left(\frac{2}{p} \right) a.$$

L 19. Let n be an integer of the form $a^2 + b^2$, where a and b are relatively prime integers and such that if p is a prime, $p \leq \sqrt{n}$, then p divides ab . Determine all such n .

L 20. If an integer n is such that $7n$ is the form $a^2 + 3b^2$, prove that n is also of that form.

L 21. Let A be the set of positive integers represented by the form $a^2 + 2b^2$, where a and b are integers and $b \neq 0$. Show that if p is a prime number and $p^2 \in A$, then $p \in A$.

L 22. Show that an integer can be expressed as the difference of two squares if and only if it is not of the form $4k + 2$ ($k \in \mathbf{Z}$).

L 23. Show that there are infinitely many positive integers which cannot be expressed as the sum of squares.

L 24. Show that any integer can be expressed as the form $a^2 + b^2 - c^2$, where $a, b, c \in \mathbf{Z}$.

L 25. Let a and b be positive integers with $\gcd(a, b) = 1$. Show that every integer greater than $ab - a - b$ can be expressed in the form $ax + by$, where $x, y \in \mathbf{N}_0$.

L 26. Let a, b and c be positive integers, no two of which have a common divisor greater than 1. Show that $2abc - ab - bc - ca$ is the largest integer which cannot be expressed in the form $xbc + yca + zab$, where $x, y, z \in \mathbf{N}_0$.

L 27. Determine, with proof, the largest number which is the product of positive integers whose sum is 1976.

L 28. (Zeckendorf) Any positive integer can be represented as a sum of Fibonacci numbers, no two of which are consecutive.

L 29. Show that the set of positive integers which cannot be represented as a sum of distinct perfect squares is finite.

L 30. Let a_1, a_2, a_3, \dots be an increasing sequence of nonnegative integers such that every nonnegative integer can be expressed uniquely in the form $a_i + 2a_j + 4a_k$, where i, j , and k are not necessarily distinct. Determine a_{1998} .

L 31. A finite sequence of integers a_0, a_1, \dots, a_n is called quadratic if for each $i \in \{1, 2, \dots, n\}$ we have the equality $|a_i - a_{i-1}| = i^2$.

(a) Prove that for any two integers b and c , there exists a natural number n and a quadratic sequence with $a_0 = b$ and $a_n = c$.

(b) Find the smallest natural number n for which there exists a quadratic sequence with $a_0 = 0$ and $a_n = 1996$.

L 32. A composite positive integer is a product ab with a and b not necessarily distinct integers in $\{2, 3, 4, \dots\}$. Show that every composite positive integer is expressible as $xy + xz + yz + 1$, with x, y, z positive integers.

L 33. Let a_1, a_2, \dots, a_k be relatively prime positive integers. Determine the largest integer which cannot be expressed in the form

$$x_1 a_2 a_3 \cdots a_k + x_2 a_1 a_3 \cdots a_k + \cdots + x_k a_1 a_2 \cdots a_{k-1}$$

for some nonnegative integers x_1, x_2, \dots, x_k .

L 34. If n is a positive integer which can be expressed in the form $n = a^2 + b^2 + c^2$, where a, b, c are positive integers, prove that, for each positive integer k , n^{2k} can be expressed in the form $A^2 + B^2 + C^2$, where A, B, C are positive integers.

L 35. Prove that every positive integer which is not a member of the infinite set below is equal to the sum of two or more distinct numbers of the set

$$\{3, -2, 2^2 3, -2^3, \dots, 2^{2k} 3, -2^{2k+1}, \dots\} = \{3, -2, 12, -8, 48, -32, 192, \dots\}.$$

L 36. Let k and s be odd positive integers such that

$$\sqrt{3k-2} - 1 \leq s \leq \sqrt{4k}.$$

Show that there are nonnegative integers t, u, v , and w such that

$$k = t^2 + u^2 + v^2 + w^2, \quad \text{and} \quad s = t + u + v + w.$$

L 37. Let $S_n = \{1, n, n^2, n^3, \dots\}$, where n is an integer greater than 1. Find the smallest number $k = k(n)$ such that there is a number which may be expressed as a sum of k (possibly repeated) elements in S_n in more than one way. (Rearrangements are considered the same.)

L 38. Find the smallest possible n for which there exist integers x_1, x_2, \dots, x_n such that each integer between 1000 and 2000 (inclusive) can be written as the sum without repetition, of one or more of the integers x_1, x_2, \dots, x_n .

L 39. In how many ways can 2^n be expressed as the sum of four squares of natural numbers?

L 40. Show that

(a) infinitely many perfect squares are a sum of a perfect square and a prime number, and

(b) infinitely many perfect squares are not a sum of a perfect square and a prime number.

L 41. *The infamous conjecture of Goldbach is the assertion that every even integer greater than 2 is the sum of two primes. Except 2, 4, and 6, every even integer is a sum of two positive composite integers : $n = 4 + (n - 4)$. What is the largest positive even integer that is not a sum of two odd composite integers?*

L 42. *Prove that for each positive integer K there exist infinitely many even positive integers which can be written in more than K ways as the sum of two odd primes.*

L 43. *A positive integer n is abundant if the sum of its proper divisors exceed n . Show that every integer greater than 89×315 is the sum of two abundant numbers.*

15. THE GEOMETRY OF NUMBERS

Srinivasa Aiyangar Ramanujan said "An equation means nothing to me unless it expresses a thought of God."

M 1. Does there exist a convex pentagon, all of whose vertices are lattice points in the plane, with no lattice point¹⁸ in the interior?

M 2. Show there do not exist four points in the Euclidean plane such that the pairwise distances between the points are all odd integers.

M 3. Prove no three lattice points in the plane form an equilateral triangle.

M 4. The lengths of a polygon with 1994 sides are $a_i = \sqrt{i^2 + 4}$ ($i = 1, 2, \dots, 1994$). Prove that its vertices are not all on lattice points.

M 5. A triangle has lattice points as vertices and contains no other lattice points. Prove that its area is $\frac{1}{2}$.

M 6. Let R be a convex region symmetrical about the origin with area greater than 4. Show that R must contain a lattice point different from the origin.

M 7. Show that the number $r(n)$ of representations of n as a sum of two squares has average value π , that is

$$\lim_{n \rightarrow \infty} \frac{1}{n} \sum_{m=1}^n r(m) = \pi.$$

M 8. Prove that on a coordinate plane it is impossible to draw a closed broken line such that

- (i) coordinates of each vertex are rational,
- (ii) the length of its every edge is equal to 1, and
- (iii) the line has an odd number of vertices.

M 9. Prove that if a lattice parallelogram contains an odd number of lattice points, then its centroid.

M 10. Prove that if a lattice triangle has no lattice points on the its boundary in addition to its vertices, and one point in its interior, then this interior point is its center of gravity.

M 11. Prove that if a lattice parallelogram contains at most three lattice points in addition to its vertices, then those are on one of the diagonals.

M 12. Find coordinates of a set of eight non-collinear planar points so that each has an integral distance from others.

¹⁸A point with integral coordinates

16. MISCELLANEOUS PROBLEMS

Mathematics is not yet ready for such problems. Paul Erdős

N 1. (a) Two positive integers are chosen. The sum is revealed to logician A , and the sum of squares is revealed to logician B . Both A and B are given this information and the information contained in this sentence. The conversation between A and B goes as follows : B starts

B : " I can't tell what the two numbers are."
 A : " I can't tell what the two numbers are."
 B : " I can't tell what the two numbers are."
 A : " I can't tell what the two numbers are."
 B : " I can't tell what the two numbers are."
 A : " I can't tell what the two numbers are."
 B : " Now I can tell what the two numbers are."

What are the two numbers ?

(b) When B first says that he cannot tell what the two numbers are, A receives a large amount of information. But when A first says that he cannot tell what the two numbers are, B already knows that A cannot tell what the two numbers are. What good does it do B to listen to A ?

N 2. It is given that 2^{333} is a 101-digit number whose first digit is 1. How many of the numbers 2^k , $1 \leq k \leq 332$, have first digit 4?

N 3. Is there a power of 2 such that it is possible to rearrange the digits giving another power of 2 ?

N 4. If x is a real number such that $x^2 - x$ is an integer, and for some $n \geq 3$, $x^n - x$ is also an integer, prove that x is an integer.

N 5. (Tran Nam Dung) Suppose that both $x^3 - x$ and $x^4 - x$ are integers for some real number x . Show that x is an integer.

N 6. Suppose that x and y are complex numbers such that

$$\frac{x^n - y^n}{x - y}$$

are integers for some four consecutive positive integers n . Prove that it is an integer for all positive integers n .

N 7. Let n be a positive integer. Show that

$$\sum_{k=1}^n \tan^2 \frac{k\pi}{2n+1}$$

is an odd integer.

N 8. The set $S = \{\frac{1}{n} \mid n \in \mathbf{N}\}$ contains arithmetic progressions of various lengths. For instance, $\frac{1}{20}, \frac{1}{8}, \frac{1}{5}$ is such a progression of length 3 and common difference $\frac{3}{40}$. Moreover, this is a maximal progression in S since it cannot be extended to the left or the right within S ($\frac{11}{40}$ and $\frac{-1}{40}$ not being members of S). Prove that for all $n \in \mathbf{N}$, there exists a maximal arithmetic progression of length n in S .

N 9. Suppose that

$$\prod_{n=1}^{1996} (1 + nx^{3^n}) = 1 + a_1x^{k_1} + a_2x^{k_2} + \cdots + a_mx^{k_m}$$

where a_1, a_2, \dots, a_m are nonzero and $k_1 < k_2 < \cdots < k_m$. Find a_{1996} .

N 10. Let p be an odd prime. Show that there is at most one non-degenerate integer triangle with perimeter $4p$ and integer area. Characterize those primes for which such triangle exist.

N 11. For each positive integer n , prove that there are two consecutive positive integers each of which is the product of n positive integers > 1 .

N 12. Let

$$\begin{array}{cccc} a_{1,1} & a_{1,2} & a_{1,3} & \cdots \\ a_{2,1} & a_{2,2} & a_{2,3} & \cdots \\ a_{3,1} & a_{3,2} & a_{3,3} & \cdots \\ \vdots & \vdots & \vdots & \ddots \end{array}$$

be a doubly infinite array of positive integers, and suppose each positive integer appears exactly eight times in the array. Prove that $a_{m,n} > mn$ for some pair of positive integers (m, n) .

N 13. The digital sum of a natural number n is denoted by $S(n)$. Prove that $S(8n) \geq \frac{1}{8}S(n)$ for each n .

N 14. Let p be an odd prime. Determine positive integers x and y for which $x \leq y$ and $\sqrt{2p} - \sqrt{x} - \sqrt{y}$ is nonnegative and as small as possible.

N 15. Let $\alpha(n)$ be the number of digits equal to one in the dyadic representation of a positive integer n . Prove that

- (a) the inequality $\alpha(n^2) \leq \frac{1}{2}\alpha(n)(1 + \alpha(n))$ holds,
- (b) the above inequality is an equality for infinitely many positive integers n , and
- (c) there exists a sequence $\{n_i\}$ such that $\lim_{i \rightarrow \infty} \frac{\alpha(n_i^2)}{\alpha(n_i)} = 0$.

N 16. Show that if a and b are positive integers, then

$$\left(a + \frac{1}{2}\right)^n + \left(b + \frac{1}{2}\right)^n$$

is an integer for only finitely many positive integer n .

N 17. Determine the maximum value of m^2+n^2 , where m and n are integers satisfying $m, n \in \{1, 2, \dots, 1981\}$ and $(n^2 - mn - m^2)^2 = 1$.

N 18. Denote by S the set of all primes p such that the decimal representation of $\frac{1}{p}$ has the fundamental period of divisible by 3. For every $p \in S$ such that $\frac{1}{p}$ has the fundamental period $3r$ one may write

$$\frac{1}{p} = 0.a_1a_2 \cdots a_{3r}a_1a_2 \cdots a_{3r} \cdots,$$

where $r = r(p)$; for every $p \in S$ and every integer $k \geq 1$ define $f(k, p)$ by

$$f(k, p) = a_k + a_{k+r(p)} + a_{k+2r(p)}.$$

a) Prove that S is finite.

b) Find the highest value of $f(k, p)$ for $k \geq 1$ and $p \in S$.

N 19. Determine all pairs (a, b) of real numbers such that $a[bn] = b[an]$ for all positive integer n . (Note that $[x]$ denotes the greatest integer less than or equal to x .)

N 20. Let n be a positive integer that is not a perfect cube. Define real numbers a, b, c by

$$a = n^{\frac{1}{3}}, \quad b = \frac{1}{a - [a]}, \quad c = \frac{1}{b - [b]},$$

where $[x]$ denotes the integer part of x . Prove that there are infinitely many such integers n with the property that there exist integers r, s, t , not all zero, such that $ra + sb + tc = 0$.

N 21. Find, with proof, the number of positive integers whose base- n representation consists of distinct digits with the property that, except for the leftmost digit, every digit differs by ± 1 from some digit further to the left.

N 22. The decimal expression of the natural number a consists of n digits, while that of a^3 consists of m digits. Can $n + m$ be equal to 2001?

N 23. Observe that

$$\begin{aligned} \frac{1}{1} + \frac{1}{3} &= \frac{4}{3}, & 4^2 + 3^2 &= 5^2, \\ \frac{1}{3} + \frac{1}{5} &= \frac{8}{15}, & 8^2 + 15^2 &= 17^2, \\ \frac{1}{5} + \frac{1}{7} &= \frac{12}{35}, & 12^2 + 35^2 &= 37^2. \end{aligned}$$

State and prove a generalization suggested by these examples.

N 24. (C. Cooper, R. E. Kennedy) A number n is called a Niven number, named for Ivan Niven, if it is divisible by the sum of its digits. For example, 24 is a Niven number. Show that it is not possible to have more than 20 consecutive Niven numbers.

N 25. Prove that if the number α is given by decimal $0.9999\dots$, where there are at least 100 nines, then $\sqrt{\alpha}$ also has 100 nines at the beginning.

N 26. Prove that there does not exist a natural number which, upon transfer of its initial digit to the end, is increased five, six, or eight times.

N 27. Which integers have the following property? If the final digit is deleted, the integer is divisible by the new number.

N 28. Let A be the set of 16 first positive integer. Find the least positive integer k satisfying the condition : In every k -subset of A , there exist two distinct $a, b \in A$ such that $a^2 + b^2$ is prime.

N 29. What is the rightmost nonzero digit of $1000000!$?

N 30. For how many positive integers n is

$$\left(1999 + \frac{1}{2}\right)^n + \left(2000 + \frac{1}{2}\right)^n$$

an integer ?

N 31. Is there a 3×3 magic square consisting of distinct Fibonacci numbers (both f_1 and f_2 may be used; that is two 1s are allowed)? (A magic square has the property that the eight sums along rows, columns, and the two main diagonals are all the same number.)

N 32. Alice and Bob play the following number-guessing game. Alice writes down a list of positive integers x_1, \dots, x_n , but does not reveal them to Bob, who will try to determine the numbers by asking Alice questions. Bob chooses a list of positive integers a_1, \dots, a_n and asks Alice to tell him the value of $a_1x_1 + \dots + a_nx_n$. Then Bob chooses another list of positive integers b_1, \dots, b_n and asks Alice for $b_1x_1 + \dots + b_nx_n$. Play continues in this way until Bob is able to determine Alice's numbers. How many rounds will Bob need in order to determine Alice's number ?

N 33. Four consecutive even numbers are removed from the set

$$A = \{1, 2, 3, \dots, n\}.$$

If the average of the remaining numbers is 51.5625, which four numbers were removed ?

N 34. Let S_n be the sum of the digits of 2^n . Prove or disprove that $S_{n+1} = S_n$ for some positive integer n .

N 35. Counting from the right end, what is the 2500th digit of $10000!$?

17. SOURCES

Divisibility Theory I

- A 1 (Ksk).¹⁹ <http://www-math.mit.edu/~kedlaya/papers>
- A 2. *Amer. Math. Monthly*, Problem 10622, Proposed by M. N. Deshpande
- A 3. *IMO 1988/6*
- A 4. *CRUX*, Problem 1420, Proposed by Shailesh Shirali
- A 5.
- A 6. *CRUX*, Problem 1746, Proposed by Richard K. Guy and Richard J. Nowakowski
- A 7. *1969 Eötvös-Kürschák Mathematics Competition*
- A 8. *Poland 2001*
- A 9 (IHH, pp. 211).
- A 10 (UmDz pp.13). *Unused Problem for the Balkan Mathematical Olympiad*
- A 11. *Slovenia 1995*
- A 12. *Putnam 1972*
- A 13. *IMO Long List 1985 P (RO2)*
- A 14. *IMO 2000/5*
- A 15. *Bulgaria 1998*
- A 16. *Slovenia 1994*
- A 17. *IMO 1967/4*
- A 18. *Amer. Math. Monthly*, Problem E2510, Proposed by Saul Singer
- A 19. *Japan 1999*
- A 20. *IMO Short List 1998*
- A 21.
- A 22. *IMO 1974/3*
- A 23 (GhEw pp.104).
- A 24. *Putnam 1996*
- A 25.
- A 26. *IMO 1972/3*

¹⁹See the References

- A 27.
- A 28. *Putnam 2000*
- A 29. *Amer. Math. Monthly, Problem E2623, Proposed by Ivan Niven*
- A 30.
- A 31. *Kazakhstan 1998*
- A 32. *IMO 1979/1*
- A 33. *IMO Short List 1996*
- A 34. *IMO Short List 2002 N3*
- A 35. *IMO Short List 2001 N4*
- A 36. *Australia 2002*
- A 37. *Bosnia and Herzegovina 2002*
- A 38. *Math. Magazine, Problem 1438, Proposed by David M. Bloom*
- A 39.
- A 40 (PJ pp.110). *UC Berkeley Preliminary Exam 1990*
- A 41 (Ae pp.137).
- A 42.
- A 43. *Iran 1994*
- A 44. *Germany 1982*
- A 45. *IMO Short List 1997*
- A 46. *Romania 1995, Proposed by I. Cucurezeanu*
- A 47 (Imv, pp. 15).
- A 48 (Imv, pp. 15).
- A 49. *IMO Short List 2001 N1*
- A 50. *Germany 2000*
- A 51. *IMO 1984/6*
- A 52. *IMO 1986/1*
- A 53. *Iran 2001*
- A 54. *IMO ShortList 1993 IND5*
- A 55. *Czech and Slovak Mathematical Olympiad 1999*
- A 56. *Romania 1987, Proposed by L. Panaitopol*
- A 57. *Bulgaria 1995*

- A 58. *APMO 2003/3*
- A 59 (AaJc, pp. 250).
- A 60. *Putnam 1988/B6*
- A 61. *Bulgaria 1995 - Arne Smeets : 2003/12/13*
- A 62. *Germany 1997 - Arne Smeets : 2003/12/13*
- A 63 (Tt). *Tournament of the Towns 2002 Fall/A-Level*
- A 64 (Tt). *Tournament of the Towns 2002 Spring/O-Level*
- A 65 (Tt). *Tournament of the Towns 2001 Fall/O-Level*
- A 66 (PeJs, pp. 5).
- A 67 (PeJs, pp. 25).
- A 68 (Her, pp. 14).
- A 69. *Math. Magazine, June 1986, Problem 1220, Proposed by Gregg Partuno*
- A 70. *Math. Magazine, Sep. 1984, Problem 1175*

Divisibility Theory II

- B 1. *IMO 1990/3 (ROM5)*
- B 2. *IMO 1999/4*
- B 3. *Ha Duy Hung : 2003/09/14*²⁰
- B 4. *APMO 1997/2*
- B 5. *APMC 2002*
- B 6. *IMO 1992/1*
- B 7. *Russia 2001*
- B 8. *IMO 1994/4*
- B 9. *IMO 2003/2*
- B 10. *IMO 2002/3*
- B 11. *IMO Short List 2000 N4*
- B 12. *Amer. Math. Monthly, Problem 10382, Proposed by Richard K. Guy*
- B 13 (Tma pp. 73).
- B 14. *India 1989*
- B 15 (ElCr pp. 11).

²⁰Contributor, Date

- B 16. *Romania 1998*
- B 17.
- B 18.
- B 19. *Turkey 1994*
- B 20. *Mediterranean Mathematics Competition 2002*
- B 21. *IMO 1998/4*
- B 22. *IMO 1977/5*
- B 23. *APMO 1998*
- B 24.
- B 25. *Iran 1998*
- B 26. *Ireland 1998*
- B 27. *Iran 1999*
- B 28. *Belarus 1999, Proposed by I. Voronovich*
- B 29. *IMO 2002/4*
- B 30. *Singapore 1997*
- B 31. *Belarus 1998, Proposed by E. Barabanov, I. Voronovich*
- B 32. *IMO 1960/1*
- B 33. *IMO 1975/4*
- B 34. *IMO Short List 1994 N7*
- B 35. *IMO Long List 1985 (TR5)*
- B 36. *IMO Long List 1987*
- B 37. *IMO Short List 1986 P10 (NL1)*
- B 38. *British Mathematical Olympiad 2003, 2-1*
- B 39. *APMO 2002/2*
- B 40. *Putnam 1997/B3*
- B 41. *Spain 1993*
- B 42. *Germany 2003 - Arne Smeets : 2003/12/13*
- B 43. *Russia 1997 - Arne Smeets : 2003/12/13*
- B 44 (EbMk, pp. 16).
- B 45 (Ams, pp. 102).
- B 46 (JDS, pp. 27).

B 47 (Rh2, pp. 98).

B 48. *Math. Magazine, Jan. 1981, Problem 1089, Proposed by M. S. Klamkin*

Arithmetic in Z_n

1. Primitive Roots

C 1.

C 2 (Imv, pp. 90).

C 3. *Math. Magazine, Problem 1419, Proposed by William P. Wardlaw*

C 4 (Km, Problems Sheet 3-9).

C 5 (Km, Problems Sheet 3-11).

C 6 (AaJc, pp. 178).

C 7 (AaJc, pp. 181).

C 8 (Her, pp. 99).

2. Quadratic Residues

C 9. *CRUX, Problem 2344, Proposed by Murali Vajapeyam*

C 10. *IMO 1996/4*

C 11 (IHH pp.147).

C 12 (Imv, pp. 72).

C 13. *Putnam 1991/B5*

C 14 (Ab, pp. 34).

3. Congruences

C 15 (Tma, pp. 127).

C 16. *Putnam 1991/B4*

C 17.

C 18. *Math. Magazine, Problem 1494, Proposed by Emeric Deutsch and Ira M. Gessel*

C 19. *Putnam 1997/B5*

C 20. *USA 1991*

C 21 (DZ). *The problem was posed by Don Zagier at the St Andrews Colloquium 1996. - Arne Smeets : 2003/12/13*

C 22. *Math. Magazine, Problem Q889, Proposed by Michael McGeachie and Stan Wagon*

C 23.

C 24.

C 25. *APMO 1991/4*

C 26 (AaJc, pp. 139).

C 27. *Putnam 1986/B3*

C 28. *Purdue POW, Spring 2003 Series/5*

C 29. *Turkey 1993*

C 30. *IMO Short List 2000 N1*

C 31. *Amer. Math. Monthly, Problem ???, Proposed by M. S. Klamkin and A. Liu*

C 32. *Turkey 2000*

C 33. *Turkey 1995*

C 34. *Belarus 1994*

C 35. *Canada 2003 - Arne Smeets : 2003/12/12*²¹

C 36. *China 1981*

C 37.

Primes and Composite Numbers

1. Composite Numbers

D 1 (DfAk, pp. 50). *Leningrad Mathematical Olympiad 1991, Here is the hint : $2z^2 = 3xy \implies x^3 + y^3 + z^3 = x^3 + y^3 + (-z)^3 - 3xy(-z)$!! In fact, It's known that $512^3 + 675^3 + 720^3 = 229 \cdot 720 \cdot 7621$.*

D 2. *IMO 2001/6*

D 3. *Math. Magazine, Problem Q614, Proposed by Rod Cooper*

D 4.

D 5. *IMO Short List 1992 P16*

D 6. *Math. Magazine, Problem Q684, Proposed by Noam Elkies*

D 7. *USA 1982*

D 8 (VsAs). - *Tran Nam Dung : 2003/12/13*²²

D 9. *IMO Short List 1996 N1*

D 10 (DfAk, pp. 9). *Leningrad Mathematical Olympiad 1987*

²¹Contributor, Date

²²Contributor, Date

- D 11 (JDS, pp. 26).
2. *Prime Numbers*
- D 12.
- D 13. *IMO Short List 1990 USS1*
- D 14 (Tma, pp. 101).
- D 15. *Math. Magazine, Problem Q789, Proposed by Norman Schaumberger*
- D 16. *Math. Magazine, Problem 1392, Proposed by George Andrews*
- D 17 (AaJc pp.212).
- D 18.
- D 19 (AaJc pp.176).
- D 20 (GjJj pp.140).
- D 21 (Ns pp.176).
- D 22. *IMO 2003/6*
- D 23. *Yugoslavia 2001*
- D 24.
- D 25 (Er pp.10). *Eötvös Competition 1896*
- D 26. *Math. Magazine, Problem 1404, Proposed by H. Gauchmen and I. Rosenholtz*
- D 27. *IMO 1989/5*
- D 28 (GjJj pp.36).
- D 29 (Tma, pp. 102).
- D 30 (Tma, pp. 128).
- D 31. *CRUX, Problem 1608, Proposed by Seung-Jin Bang*
- D 32. *Singapore 1989*
- D 33. *Russia 1996*
- D 34 (Rh, pp. 43).
- D 35 (Tt). *Tournament of the Towns 2001 Fall/O-Level*
- D 36 (PeJs, pp. 160).
- D 37 (PeJs, pp. 174).
- D 38 (PbAw, pp. 1).
- D 39 (Wsa, pp. 217).

D 40 (DNI, 19).

D 41 (Her, pp. 11).

Rational and Irrational Numbers

Rational Numbers

E 1 (Vvp, pp. 40).

E 2. *CRUX, Problem 1632, Proposed by Stanley Rabinowitz*

E 3. *IMO ShortList 1991 P19 (IRE 5)*

E 4. *IMO Long List 1967 P20 (DDR)*

E 5. *Austria 2002*

E 6. *Romania 2001, Proposed by Marius Ghergu*

E 7. *IMO Long List 1967 (GB)*

E 8. *Poland 2002*

E 9. *IMO Short List 1999*

E 10. *Berkeley Math Circle Monthly Contest 1999-2000*

E 11. *Iran 1998*

E 12 (Zh, PP. 40). *Mediterranean Mathematics Competition 1999 (Proposed by Ukraine)*

E 13 (PeJs, pp. 99).

E 14 (PbAw, pp. 2).

E 15 (EbMk, pp. 15).

E 16 (EbMk, pp. 23).

E 17 (GML, pp. 167).

E 18 (GML, pp. 183).

Irrational Numbers

E 19. *The Grosman Memorial Mathematical Olympiad 1999*

E 20. *Belarus 2002*

E 21. *Belarus 2001*

E 22. *Putnam 1980*

E 23. *CRUX, Problem A240, Proposed by Mohammed Aassila*

E 24.

E 25.

E 26.

E 27.

E 28.

E 29.

E 30. *CRUX, Problem 2305, Proposed by Richard I. Hess*

E 31. *Amer. Math. Monthly, Problem 10630, Proposed by Richard Strong*

E 32. *Math. Magazine, Problem 1372, Proposed by Nick Lord*

E 33 (AI, pp. 106). *For a proof, See [Kmh].*

E 34 (KhKw, pp. 11).

E 35 (AI, pp. 116). *For a proof, See [KaMr].*

E 36. *USA 1973*

E 37. *IMO 1987/5*

E 38. *APMO 1994/5*

E 39.

E 40. *Putnam 1995*

E 41. *Math. Magazine, Problem 1385, Proposed by Howard Morris*

E 42 (Ae, pp. 226).

E 43 (PeJs, pp. 95).

E 44 (PeJs, pp. 99).

E 45 (PbAw, pp. 2).

E 46 (Ams, pp. 14).

Diophantine Equations I

F 1. *AIME 1989/9*

F 2. *UC Berkeley Preliminary Exam 1983*

F 3. *Taiwan 1998*

F 4.

F 5.

F 6.

F 7. *Poland 2002*

F 8. *Bulgaria 1999*

F 9. *Ireland 1995*

- F 10. *Putnam 2001*
- F 11. *India 1998*
- F 12.
- F 13. *Italy 1994*
- F 14. *Canada 1991*
- F 15. *Balkan Mathematical Olympiad 1998*
- F 16. *Poland 2003*
- F 17. *Vietnam 2002*
- F 18. *Romania 1995, Proposed by M. Becheanu*
- F 19 (UmDz pp.14). *Unused Problem for the Balkan Mathematical Olympiad*
- F 20. *APMO 1993/4*
- F 21. *APMO 1989/2*
- F 22. *Poland 1998*
- F 23.
- F 24. *IMO 1982/4*
- F 25. *IMO Short List 2002 N1*
- F 26. *Ukraine 2002*
- F 27. *IMO Short List 2000 N5*
- F 28. *IMO Short List 1997 N6*
- F 29. *Belarus 2000*
- F 30. *IMO Short List 1993 GEO3*
- F 31 (Eb1, pp. 19). *Amer. Math. Monthly 61(1954), 126; 62(1955), 263*
- F 32. *IMO Long List 1987 (Romaina)*
- F 33. *IMO Long List 1967 P (PL)*
- F 34. *IMO Long List 1985 (SE1)*
- F 35. *IMO Long List 1985 (TR3)*
- F 36. *Italy 1996*
- F 37. *Bulgaria 1996 - Arne Smeets : 2003/12/12*
- F 38 (JeMm, pp. 10).
- F 39 (KhKw, pp. 9).
- F 40 (EbMk, pp. 44).

- F 41 (Wsa, pp. 39).
 F 42 (GML, pp. 2).
 F 43 (GML, pp. 33).
 F 44 (GML, pp. 192).
 F 45 (Ljm, pp. 21).
 F 46 (Ljm, pp. 57).
 F 47 (Ljm, pp. 94).
 F 48 (Ljm, pp. 205).
 F 49 (Ljm, pp. 208).
 F 50 (Her, pp. 287).
 F 51 (Rh3, pp. 207).
 F 52. *Tom Verhoeff : 2004/05/23*²³
 F 53. *Math. Magazine, Jan. 1985, Lorraine L. Foster, pp. 50-51*
 F 54. *Math. Magazine, Sep. 1980, Problem 1077, Proposed by Henry Klostergaard*

Diophantine Equations II

- G 1. *British Mathematical Olympiad 2002/2003, 1-1*
 G 2. *Latvia 1995*
 G 3 (MaGz pp.13-16).
 G 4. *Belarus 1999, Proposed by S. Shikh*
 G 5. *IMO Long List 1987 (Greece)*
 G 6 (VsAs). - *Tran Nam Dung : 2003/12/13*
 G 7. *CRUX, Problem 1797, Proposed by Marcin E. Kuczma*
 G 8. *India 1995*
 G 9.
 G 10 (Rdc pp.51).
 G 11. *Austria 2002*
 G 12. *Austria 2002*
 G 13. *IMO Short List 2002 N4*
 G 14. *IMO Short List 2001 N2*

²³Contributor, Date

- G 15. *Hong Kong 2000*
- G 16. *Belarus 1999*
- G 17. *Baltic Way 2002*
- G 18.
- G 19. *IMO 1997/5*
- G 20. *Belarus 2000*
- G 21. *Iran 1998*
- G 22. *Belarus 1996*
- G 23. *Korea 1998*
- G 24. *Hungary 1998*
- G 25 (Eb2, pp. 20). *Q657, Math. Magazine 52(1979), 47, 55*
- G 26. *IMO Short List 1980 (GB)*
- G 27 (DfAk, pp. 18). *Leningrad Mathematical Olympiad 1988*
- G 28. *British Mathematical Olympiad 2002/2003, 1-5*
- G 29. *British Mathematical Olympiad 2000, 2-3*
- G 30. *Math. Magazine, Problem 1529, Proposed by David C. Kay*
- G 31. *Math. Magazine, Problem 1538, Proposed by Murray S. Klamkin and George T. Gilbert.*
- G 32. *CRUX, Problem 2351, Proposed by Paul Yiu*
- G 38 (EbMk, pp. 22).
- G 33 (EbMk, pp. 46).
- G 34 (DNI, 42).
- G 35. *Vietnam 2004*
- G 36 (JDS, pp. 29).

Functions in Number Theory

1. Floor Function and Fractional Part Function

- H 1. *IMO ShortList 1991 P20 (IRE 3)*
- H 2 (EbMk, pp. 5).
- H 3 (EbMk, pp. 7).
- H 4. *Canada 1987*
- H 5. *CRUX, Problem 1650, Proposed by Iliya Bluskov*

H 6. *Iran 1996*

H 7. *Math. Magazine, Problem 1410, Proposed by Seung-Jin Bang*

H 8. *Canadian Mathematical Society Notes, Problem P11, Proposed by Mihály Bencze*

H 9. *Taiwan 1998*

H 10. *Amer. Math. Monthly, Problem 10346, Proposed by David Doster*

H 11. *Korea 2000*

H 12 (IHH pp.142).

H 13. *CRUX, Problem 2321, Proposed by David Doster*

H 14.

H 15. *APMO 1993/2*

H 16.

H 17. *Bulgaria 2003 - Arne Smeets : 2003/12/13*

H 18 (Tt). *Tournament of the Towns 2002 Spring/A-Level*

H 19 (PbAw, pp. 5).

H 20 (Ams, pp. 45).

2. Euler phi Function

H 21.

H 22.

H 23. *Math. Magazine, Problem 1376, Proposed by Eric Canning*

H 24. *Amer. Math. Monthly, Problem 10837, Proposed by Hojoo Lee*

H 25 (Km, Problems Sheet 1-11).

H 26. *Amer. Math. Monthly, Problem 10626, Proposed by Florian Luca*

H 27 (Rdc pp.36).

H 28 (PeJs, pp. 237).

H 29 (Pr, pp. 38).

H 30. *Math. Magazine, Sep. 1983, Problem 1153, Proposed by Charles R. Wall*

3. Divisor Functions

H 31. *Russia 1998*

H 32. *Canada 1999*

H 33. *IMO 1998/3*

H 34. *IMO Short List 2000 N2*

H 35. *Canada 1992*

H 36. *IMO Short List 1983 (Belgium)*

H 37 (Rh pp.104). *Quantum, Problem M59, Contributed by B. Martynov*

H 38 (PeJs, pp. 237).

H 39 (PeJs, pp. 237).

H 40 (Jjt, pp. 95).

H 41 (Jjt, pp. 169).

H 42 (Dmb, pp. 108).

4. More Functions

H 43 (Tau). *and [Tau2]*

H 44. *Germany 1996*

H 45. *Belarus 1999, Proposed by D. Bazylev*

H 46 (Tma, pp.175).

H 47.

5. Functional Equations

H 48. *Singapore 1996*

H 49. *Turkey 1995*

H 50. *IMO 1977/6*

H 51. *Unknown*²⁴

H 52. *Unknown*

H 53. *IMO unused 1997*

H 54. *Balkan 2002*

H 55. *USAMO Summer Program 2001*

H 56. *Austria 1989*

H 57. *IMO 1996/3*

H 58. *Canada 2002*

H 59. *Canada 1969*

H 60. *Slovenia 1997*

²⁴I don't know the origin of the problem.

- H 61. *Unknown*
- H 62. *APMC 1997*
- H 63. *South Africa 1997*
- H 64. *Belarus 1999*
- H 65. *APMC 1984*
- H 66. *Unknown*
- H 67. *Nordic Mathematics Contest 1998*
- H 68. *KMO Winter Program Test 2001*
- H 69. *Ukraine 1997*
- H 70. *IMO 1990/4*
- H 71. *IMO 1988/3*
- H 72. *IMO 1998/6*
- H 73. *IMO 1982/1*
- H 74. *IMO Short List 1988*
- H 75. *Romania 1986*
- H 76. *Iran 1995*
- H 77.
- H 78.
- H 79.
- H 80.
- H 81.
- H 82.

Polynomials

- I 1. *Math. Magazine, Problem Q800, Proposed by Bjorn Poonen*
- I 2.
- I 3. *IMO 1987/6*
- I 4. *Balkan Mathematical Olympiad 1989*
- I 5 (Twh, pp. 111).
- I 6 (Twh, pp. 114).
- I 7. *IMO 1993/1*
- I 8 (Ac, pp. 87). *For a proof, see [En].*

I 9. *CRUX, Problem A230, Proposed by Naoki Sato*

I 10 (Ae, pp. 257).

I 11 (Ae, pp. 258).

I 12 (DNI, 47).

I 13. *Math. Magazine, Jan. 1982, Problem 1113, Proposed by William H. Gustafson*

Sequences of Integers

1. Linear Recurrences

J 1. *IMO Short List 1988*

J 2 (Nv pp.58).

J 3 (Nv pp.74).

J 4 (Nv pp.75).

J 5 (Eb1 pp.21).

J 6. *Math. Magazine, Problem 1390, Proposed by J. F. Stephany*

J 7. *Canada 1998*

J 8. *Vietnam 1999*

J 9. *Germany 1995 - Arne Smeets : 2003/11/24*

J 10. *Bulgaria 2003 - Arne Smeets : 2003/12/13*

J 11. *Math. Magazine, Problem 1558, Proposed by Mansur Boase*

J 12 (Ae, pp. 226).

J 13 (Rh2, pp. 197).

2. Recursive Sequences

J 14.

J 15. *Amer. Math. Monthly, Problem E2619, Proposed by Thomas C. Brown*

J 16. *Putnam 1983*

J 17. *Putnam 1999*

J 18. *United Kingdom 1998*

J 19. *Serbia 1998*

J 20. *United Kingdom 2002*

J 21 (KiMr pp. 233).

J 22. *Putnam 1983*

- J 23.** *APMC 1999*
- J 24.** *IMO Short List 2001 N3*
- J 25.** *Poland 2002*
- J 26** (Ae pp.228).
- J 27.** *IMO Short List 1994 N6*
- J 28.** *IMO Short List 1990 HUN1*
- J 29.** *Putnam 1985/A4*
- J 30.** *CRUX, Problem 2446, Proposed by Catherine Shevlin*
- J 31.** *Netherlands 1994 - Arne Smeets : 2003/12/12*
- J 32** (Tt). *Tournament of the Towns 2002 Fall/A-Level- Arne Smeets : 2003/12/12*
- J 33** (Tt). *Tournament of the Towns 2003 Spring/O-Level*
- J 34** (JtPt, pp. 93). *Russia 1984 - Arne Smeets : 2003/12/12*
- J 35.** *USA 1993*
- J 36.** *IMO Long List 1987 (GB)*
- J 37.** *Vietnam 2000 - Tran Nam Dung : 2003/12/13*
- J 38.** *Vietnam 1999 - Tran Nam Dung : 2003/12/13*
- J 39.** *Poland 1995 - Arne Smeets : 2003/12/13*
- J 40.** *Canada 1986 - Arne Smeets : 2003/12/13*
- J 41.** *China 1991*
- J 42.** *Math. Magazine, Problem 1545, Proposed by Erwin Just*
- J 43** (Rh, pp. 276).
- J 44** (Tt). *Tournament of the Towns 2002 Fall/O-Level*
- J 45** (Tt). *Tournament of the Towns 2002 Spring/A-Level*
- J 46** (Ae, pp. 227).
- J 47** (Ams, pp. 104).
- J 48** (Rh3, pp. 103).
3. *More Sequences*
- J 49.** *IMO Long List 1985 (RO3)*
- J 50.** *IMO Short List 1991 P14 (USS 2)*
- J 51.** *IMO 1991/2*

- J 52.** *IMO Short List 1993*
- J 53.** *IMO Short List 1999 N3*
- J 54.** *Pi Mu Epsilon Journal, Problem 339, Proposed by Paul Erdős*
- J 55.** *Iran 1998*
- J 56.** *IMO Short List 1989*
- J 57.** *Taiwan 1996*
- J 58.** *USA 2002*
- J 59.** *Putnam 1993/A6*
- J 60.** *Vietnam 2001 - Tran Nam Dung : 2003/12/13*
- J 61.** *Croatia 1994*
- J 62 (Tt).** *Tournament of the Towns 2003 Spring/O-Level*
- J 63 (Tt).** *Tournament of the Towns 2001 Fall/A-Level*
- J 64 (Tt).** *Tournament of the Towns 2001 Fall/A-Level*
- J 65 (GML,** pp. 173).
- Combinatorial Number Theory***
- K 1 (Rh2,** pp. 243).
- K 2.** *IMO Long List 1985 (PL2)*
- K 3.** *IMO 1971/3*
- K 4.** *Berkeley Math Circle Monthly Contest 1999-2000*
- K 5.** *India 1998*
- K 6.** *USA 2001*
- K 7.** *USA 1998*
- K 8.** *Romania 1998*
- K 9.** *India 1998*
- K 10.** *Romania 1998*
- K 11.** *IMO 1991/3*
- K 12.** *Math. Magazine, Problem 1466, Proposed by David M. Bloom*
- K 13.** *IMO 1985/2*
- K 14.** *Romania 1997, Proposed by Marian Andronache and Ion Savu*
- K 15.** *IMO 1983/5*
- K 16.** *IMO Short List 2001*

- K 17. *Turkey 1996*
- K 18. *IMO 1995/6*
- K 19. *IMO Short List 2002 N5*
- K 20. *IMO Short List 1998 P16*
- K 21. *IMO Short List 1998 P17*
- K 22. *IMO ShortList 1998 P20*
- K 23. *IMO Short List 1996*
- K 24 (TaZf pp.10). *High-School Mathematics (China) 1994/1*
- K 25. *IMO Short List 2002 A6*
- K 26. *IMO Short List 2001 C4*
- K 27. *IMO Short List 2000 C6*
- K 28. *IMO Short List 1999 A4*
- K 29. *IMO Short List 1999 C4*
- K 30. *Australia 2002*
- K 31 (Ae pp.228).
- K 32. *IMO ShortList 1991 P24 (IND 2)*
- K 33. *IMO Short List 1990 CZE3*
- K 34. *IMO Short List 1990 MEX2*
- K 35. *USA 2002*
- K 36. *IMO Short List 1995*
- K 37. *IMO Short List 1999*
- K 38. *IMO Short List 1999*
- K 39. *Romania 1995*
- K 40. *Japan 1990*
- K 41. *USA 2003*
- K 42. *Putnam 1997/A5*
- K 43. *British Mathematical Olympiad 2000 - Arne Smeets : 2003/12/13*
- K 44. *Bulgaria 2003 - Arne Smeets : 2003/12/13*
- K 45. *IMO 1970/4*
- K 46. *CRUX, Problem A233, Proposed by Mohammed Aassila*
- K 47 (KhKw, pp. 12).

- K 48** (KhKw, pp. 13).
- K 49** (Ams, pp. 12).
- K 50** (Prh, pp. 29).
- K 51** (DNI, 19).
- K 52** (JDS, pp. 31).
- K 53** (Her, pp. 16).
- K 54** (Rh2, pp. 89).
- K 55** (Rh2, pp. 125).
- K 56** (Rh2, pp. 145).
- K 57** (Rh2, pp. 202).
- K 58** (Rh3, pp. 213).
- K 59** (Rh3, pp. 51).

Additive Number Theory

- L 1.** *Amer. Math. Monthly, Problem 10426, Proposed by Noam Elkies and Irving Kaplanky*
- L 2.** *Netherlands 1994 - Arne Smeets : 2003/12/12*
- L 3.** *Belarus 2002 Proposed by V. Bernik - Arne Smeets : 2003/12/13*
- L 4.** *Math. Magazine, Problem Q817, Proposed by Robert B. McNeill*
- L 5.**
- L 6** (IHH, pp. 474).
- L 7** (Tma, pp. 22).
- L 8** (Rdc pp.24).
- L 9.** *IMO Short List 1990 AUS3*
- L 10.** *IMO 1992/6*
- L 11.** *IMO 1997/6*
- L 12** (Hua pp.199).
- L 13.** *Math. Magazine, Problem Q814, Proposed by Paul Erdős*
- L 14.** *Romania 2001, Proposed by Laurentiu Panaitopol*
- L 15.** *Amer. Math. Monthly, Problem E3064, Proposed by Ion Cucurezeanu*
- L 16.** *Putnam 2000*
- L 17.**

- L 18. *Amer. Math. Monthly, Problem 2760, Proposed by Kenneth S. Williams*
- L 19. *APMO 1994/3*
- L 20. *India 1998*
- L 21. *Romania 1997, Proposed by Marcel Tena*
- L 22.
- L 23.
- L 24.
- L 25.
- L 26. *IMO 1983/3*
- L 27. *IMO 1976/4*
- L 28.
- L 29. *IMO Short List 2000 N6*
- L 30. *IMO Short List 1998 P21*
- L 31. *IMO Short List 1996 N3*
- L 32. *Putnam 1988/B1*
- L 33. *Math. Magazine, Problem 1561, Proposed by Emre Alkan*
- L 34 (KhKw, pp. 21).
- L 35 (EbMk, pp. 46).
- L 36 (Wsa, pp. 271).
- L 37 (GML, pp. 37).
- L 38 (GML, pp. 144).
- L 39 (DNI, 28).
- L 40 (JDS, pp. 25).
- L 41 (JDS, pp. 25).
- L 42. *Math. Magazine, Feb. 1986, Problem 1207, Proposed by Barry Powell*
- L 44. *Math. Magazine, Nov. 1982, Problem 1130, Proposed by J. L. Selfridge*

The Geometry of Numbers

- M 1. *Math. Magazine, Problem 1409, Proposed by Gerald A. Heur*
- M 2. *Putnam 1993/B5*

M 3.

M 4. *Israel 1994*

M 5.

M 6 (Hua pp.535).

M 7 (GjJj pp.215).

M 8. *IMO Short List 1990 USS3*

M 9 (PeJs, pp. 125).

M 10 (PeJs, pp. 125).

M 11 (PeJs, pp. 125).

M 12 (Jjt, pp. 75).

Miscellaneous Problems

N 1. *Math. Magazine, May 1984, Problem 1173, Proposed by Thomas S. Ferguson*

N 2 (Tt). *Tournament of the Towns 2001 Fall/A-Level*

N 3 (Pt). *Tournament of the Towns*

N 4. *Ireland 1998*

N 5. *Vietnam 2003 - Tran Nam Dung : 2003/12/13*

N 6. *Amer. Math. Monthly, Problem E2998, Proposed by Clark Kimberling*

N 7.

N 8. *British Mathematical Olympiad 1997 - Arne Smeets : 2003/12/13*

N 9. *Turkey 1996 - Arne Smeets : 2003/12/12*

N 10. *CRUX, Problem 2331, Proposed by Paul Yiu*

N 11 (Rh, pp. 165). *Unused problems for 1985 Canadian Mathematical Olympiad*

N 12. *Putnam 1985/B3*

N 13. *Latvia 1995*

N 14. *IMO Short List 1992 P17*

N 15.

N 16 (Ns pp.4).

N 17. *IMO 1981/3*

N 18. *IMO Short List 1999 N4*

- N 19.** *IMO Short List 1998 P15*
- N 20.** *IMO Short List 2002 A5*
- N 21.** *USA 1990*
- N 22** (Tt). *Tournament of the Towns 2001 Spring/O-Level*
- N 23** (EbMk, pp. 10).
- N 24** (Jjt, pp. 58).
- N 25** (DNI, 20).
- N 26** (DNI, 12).
- N 27** (DNI, 11).
- N 28.** *Vietnam 2004*
- N 29** (JDS, pp. 28).
- N 30** (JDS, pp. 30).
- N 31** (JDS, pp. 31).
- N 32** (JDS, pp. 57).
- N 33** (Rh2, pp. 78).
- N 34.** *Math. Magazine, Nov. 1982, Q679, Proposed by M. S. Klamkin and M. R. Spiegel*
- N 35.** *Math. Magazine, Sep. 1980, Problem 1075, Proposed by Phillip M. Dunson*

18. REFERENCES

- AaJc Andrew Adler, John E. Coury, *The Theory of Numbers - A Text and Source Book of Problems*, John and Bartlet Publishers
- Ab Alan Baker, *A Consise Introduction to the Theory of Numbers*, Cambridge University Press
- Ac Allan Clark, *Elements of Abstract Algebra*, Dover
- Ae Arthur Engel, *Problem-Solving Strategies*, Springer-Verlag
- Ams A. M. Slinko, *USSR Mathematical Olympiads 1989-1992*, AMT²⁵
- AI A. N. Parshin, I. R. Shafarevich, *Number Theory IV - Encyclopaedia of Mathematical Sciences, Volume 44*, Spinger-Verlag
- DfAk Dmitry Fomin, Alexey Kirichenko, *Leningrad Mathematical Olympiads 1987-1991*, MathPro Press
- Dmb David M. Burton, *Elementary Number Theory*, MathPro Press
- DNI D. O. Shklarsky, N. N. Chentzov, I. M. Yaglom, *The USSR Olympiad Problem Book*, Dover
- Dz <http://www-gap.dcs.st-and.ac.uk/~john/Zagier/Problems.html>
- Eb1 Edward J. Barbeau, *Pell's Equation*, Springer-Verlag
- Eb2 Edward J. Barbeau, *Power Play*, MAA²⁶
- EbMk Edward J. Barbeau, Murry S. Klamkin *Five Hundred Mathematical Challenges*, MAA
- ElCr Edward Lozansky, Cecil Rousseau, *Winning Solutions*, Springer-Verlag
- En Eugen Netto, ??, *Mathematische Annalen*, vol 48(1897)
- Er Elvira Rapaport, *Hungarian Problem Book I*, MAA
- GhEw G. H. Hardy, E. M. Wright, *An Introduction to the theory of numbers*, Fifth Edition, Oxford University Press
- GjJj Gareth A. Jones, J. Mary Jones, *Elementary Number Theory*, Springer-Verlag
- GML George T. Gilbert, Mark I. Krusemeyer, Loren C. Larson, *The Wohas-cum County Problem Book*, MAA
- Her H. E. Rose, *A Course in Number Theory*, Cambridge University Press
- Hs *The MacTutor History of Mathematics Archive*, <http://www-gap.dcs.st-and.ac.uk/~history/index.html>
- Hua Hua Loo Keng, *Introduction to Number Theory*, Springer-Verlag
- IHH Ivan Niven, Herbert S. Zuckerman, Hugh L. Montgomery, *An Introduction to the Theory of Numbers*, Fifth Edition, John Wiley and Sons, Inc.
- Imv I. M. Vinogradov, *An Introduction to The Theory of Numbers*, Pergamon Press

²⁵Australian Mathematics Trust

²⁶Mathematical Association of America

- JDS Joseph D. E. Konhauser, Dan Velleman, Stan Wagon, *Which Way Did The Bicycle Go?*, MAA
- Jjt James J. Tattersall, *Elementary Number Theory in Nine Chapters*, Cambridge University Press
- JtPt Jordan B. Tabov, Peter J. Taylor, *Methods of Problem Solving Book 1*, AMT
- JeMm J. Esmonde, M. R. Murty, *Problems in Algebraic Number Theory*, Springer-Verlag
- KaMr K. Alladi, M. Robinson, *On certain irrational values of the logarithm*. Lect. Notes Math. 751, 1-9
- KhKw Kenneth Hardy, Kenneth S. Williams, *The Green Book of Mathematical Problems*, Dover
- KiMr Kenneth Ireland, Michael Rosen, *A Classical Introduction to Modern Number Theory*, Springer-Verlag
- Km Keith Matthews, *MP313 Number Theory III, Semester 2, 1999*, <http://www.numbertheory.org/courses/MP313/index.html>
- Kmh K. Mahler, *On the approximation of π* , Proc. Kon. Akad. Wetensch. A., Vol 56, 30-42
- Ksk Kiran S. Kedlaya, *When is $(xy+1)(yz+1)(zx+1)$ a square?*, Math. Magazine, Vol 71 (1998), 61-63
- Ljm L. J. Mordell, *Diophantine Equations*, Academic Press
- MaGz Martin Aigner, Günter M. Ziegler, *Proofs from THE BOOK*, Springer-Verlag
- Nv Nicolai N. Vorobiev, *Fibonacci Numbers*, Birkhäuser
- PbAw Pitor Biler, Alfred Witkowski, *Problems in Mathematical Analysis*, Marcel Dekker, Inc.
- PJ Paulo Ney de Souza, Jorge-Nuno Silva, *Berkeley Problems in Mathematics*, Second Edition, Springer-Verlag
- Pp *Purdue Univ. POW*, <http://www.math.purdue.edu/academics/pow>
- Pr Paulo Ribenboim, *The New Book of Prime Number Records*, Springer-Verlag
- Hrh Paul R. Halmos, *Problems for Mathematicians, Young and Old*, MAA
- Pt Peter J. Taylor, *Tournament of the Towns 1984-1989, Questions and Solutions*, AMT
- PeJs Paul Erdős, János Surányi, *Topics in the Theory of Numbers*, Springer-Verlag
- Rc Robin Chapman, *A Polynomial Taking Integer Values*, Math. Magazine, Vol 69 (1996), 121
- Rdc Robert D. Carmichael, *The Theory of Numbers*
- Rh R. Honsberger, *Mathematical Chestnuts from Around the World*, MAA
- Rh2 R. Honsberger, *In Pólya's Footsteps*, MAA
- Rh3 R. Honsberger, *From Erdős To Kiev*, MAA

- Rs Rizos Sakellariou, *On Equal Sums of Like Powers (Euler's Conjecture)*, <http://www.cs.man.ac.uk/cnc/EqualSums/equalsums.html>
- Tau *Tau Function*, <http://mathworld.wolfram.com/TauFunction.html>
- Tau2 A000594, <http://www.research.att.com/cgi-bin/access.cgi/as/njas/sequences/eisA.cgi?Anum=A000594>
- TaZf Titu Andreescu, Zuming Feng, *102 Combinatorial Problems From the Training of the USA IMO Team*, Birkhäuser
- Tma Tom M. Apostol, *Introduction to Analytic Number Theory*, Springer-Verlag
- Twh Thomas W. Hungerford, *ABSTRACT ALGEBRA - An Introduction*, Brooks/Cole
- UmDz Uroš Milutinović, Darko Žubrinić, *Balkanian Mathematical Olympiades 1984-1991*
- VsAs V. Senderov, A. Spivak, *Fermat's Little Theorem*, Quantum, May/June 2000
- Vvp V. V. Prasolov, *Problems and Theorems in Linear Algebra*, AMS²⁷
- Wlp Kiran S. Kedlaya, *William Lowell Putnam Mathematics Competition Archive*, <http://www.unl.edu/amc/a-activities/a7-problems/putnam/-pindex.html>
- Wsa W. S. Anglin, *The Queen of Mathematics*, Kluwer Academic Publishers
- Zh Zeljko Hanjcs, *Mediterranean Mathematics Competition MMC*, Mathematics Competitions, Vol 12 (1999), 37-41

²⁷American Mathematical Society